

ANNUAL INDEX

VOLUME 2 1967

AUTHORS

A

- Ackman, R. G. Letter to the editor re the chain-length overlap problem in gas-liquid chromatography with polyester liquid phases, 502
 — and R. P. Hansen. The occurrence of diastereomers of phytanic and pristanic acids and their determination by gas-liquid chromatography, 357
 — J. C. Sipos and P. M. Jangaard. A quantitative problem in the open tubular gas chromatography of fatty acid esters from cod liver lipids, 251
 Allen, E., A. R. Johnson, A. C. Fogerty, Judith A. Pearson and F. S. Shenstone. Inhibition by cyclopropene fatty acids of the desaturation of stearic acid in hen liver, 419
 Anderson, Beverly S. See Alec D. Keith
 Anderson, Robert E., Nestor R. Bottino and Raymond Reiser. Short communication re pancreatic lipase hydrolysis as a source of diglycerides for the stereospecific analysis of triglycerides, 440
 Ansell, G. B. See L. A. Horrocks
 Appleqvist, Lars-Ake, and Karl-Axel Melin. Short communication re an autosampler for solvent-free sample introduction into a gas chromatograph, 351

B

- Baer, Eric, and Ranga Robinson. Short communication re phosphonolipids. IX. Phosphonolipid metabolites: synthesis of L- α -glyceryl-(2-trimethylammoniumethyl) phosphonate, 194; erratum, 286
 Barclay, A. S. See J. M. Hagemann
 Bauman, A. J., Ray E. Cameron, Gene Kritchevsky and George Rouser. Short communication re detection of phthalate esters as contaminants of lipid extracts from soil samples stored in standard soil bags, 85; erratum, 198
 Beal, R. E. See L. T. Black
 Berrettoni, J. N. See Antanas Butkus
 Black, L. T., and R. E. Beal. Short communication re determination of unsaturation in oils in the presence of saturated aldehydes, 280
 Bollinger, James N. The isolation and tentative identification of diglycerol ethers from the Walker 256 carcinoma of the rat and a human lymphosarcoma, 143
 Bottino, Nestor R. Metabolism of unsaturated fatty acids in the intestine, 155
 — See Robert E. Anderson
 — Gloria A. Vandenberg and Raymond Reiser. Resistance of certain long-chain polyunsaturated fatty acids of marine oils to pancreatic lipase hydrolysis, 489
 Brenner, Rodolfo R. See Angel Catalá
 — See O. Mercuri
 Brown, Thomas O. See Donald R. Van Harken
 Burren, C. H., I. Ehrlich and P. Johnson. Excretion of lipids by the liver fluke (*Fasciola Hepatica* L.), 353
 Butkus, Antanas, and J. N. Berrettoni. Quantitative and qualitative lipid correlation in experimental endogenous hyperlipemia, 212
 Buziassy, Charles. See Ann C. Noble

C

- Cameron, Ray E. See A. J. Bauman
 Carpenter, D. L. See J. Sampugna
 Catalá, Angel, and Rodolfo R. Brenner. Relative incorporation of linoleic and arachidonic acid in phospholipids and triglycerides of different rat tissues, 114

- Cheng, Shu-Sing, Claude Piantadosi and Fred Snyder. Short communication re lipid droplets and glyceryl ether diesters in Ehrlich ascites cells grown in tissue culture, 193
 Chiu, Mei. See C. K. Chow
 — See H. H. Draper
 Chow, C. K., H. H. Draper, A. S. Csallany and Mei Chiu. The metabolism of C₁₄- α -tocopheryl quinone and C₁₄- α -tocopheryl hydroquinone, 390
 Christie, W. W., and F. D. Gunstone. Letter to the editor re naming diastereoisomeric polyhydroxystearates, 286
 Cole, Francis E. See Ronald C. Reitz
 Coniglio, J. G., F. B. Culp and A. Goswami. The metabolism of 1-¹⁴C arachidonic acid in pyridoxine-deficient and pair-fed control rats, 484
 Cornatzer, W. E. See Shuang-Shine Tsao
 Csallany, A. S. See C. K. Chow
 — See H. H. Draper
 Culp, F. B. See J. G. Coniglio
 Cwiaknia, Charles E. See Emanuel Merdinger

D

- Das, Manik L., and George Rouser. Lipid composition of beef heart ventricle, 1
 Davison, V. L. See R. Kleiman
 Day, E. A. See C. Jane Wyatt
 Del Rosario, R. R. See Ki Soon Rhee
 Deluca, H. F. See J. L. Moore
 Derrick, Nancy M., and Lawrence A. Wishner. Autoxidation of tissue lipids. I. Incorporation of dietary fatty acids and formation of monocarbonyl compounds in adipose tissue lipids of the vitamin E-deficient rat, 133
 Diver, M. J. See W. D. Mitchell
 Dolev, Ami. See C. D. Evans
 — W. K. Rohwedder, T. L. Mounts and H. J. Dutton. Letter to the editor re a note on the mechanism of lipoxidase reaction and the origin of the oxygen incorporated into linoleate hydroperoxide, 442
 — W. K. Rohwedder and H. J. Dutton. Mechanism of lipoxidase reaction. I. Specificity of hydroperoxidation of linoleic acid, 28
 — W. K. Rohwedder, T. L. Mounts and H. J. Dutton. Mechanism of lipoxidase reaction. II. Origin of the oxygen incorporated into linoleate hydroperoxide, 33
 Draper, H. H. See C. K. Chow
 — A. S. Csallany and Mei Chiu. Isolation of a trimer of α -tocopherol from mammalian liver, 47
 Dubuc, J. See M. Kraml
 Duffy, Thomas E., John J. O'Neill and A. N. Siakotos. Short communication re effect of phenazine methosulfate on the incorporation of ¹⁴C-labeled glucose into lipids of newborn brain, 195
 Dugan, L. R., Jr. See Ki Soon Rhee
 Dutton, H. J. See Ami Dolev
 — See R. Kleiman
 — See Howard W. Sprecher
 Dvornik, D. See M. Kraml

E

- Earle, F. R. See J. M. Hagemann
 — See R. Kleiman
 Eddy, C. R. See D. A. Lutz
 Ehrlich, I. See C. H. Burren
 Evans, C. D., G. R. List, Ami Dolev, D. G. McConnell and R. L. Hoffmann. Pentane from thermal decomposition of lipoxidase-derived products, 432

F

- Favarger, P. See S. Rous
 Fewster, Mona E., and Michael O. Hall. The renal phospholipid composition of choline-deficient rats, 239
 Fillerup, Dorothy L., and James F. Mead. The lipids of the aging human brain, 295
 Fogerty, A. C. See E. Allen
 — See A. R. Johnson
 Freeman, N. K. See F. T. Hatch

G

- Galli, Claudio, and Daniela Re Cecconi. Lipid changes in rat brain during maturation, 76
 Galli, Giovanni, and Enrica G. Paoletti. Separation of cholesterol-desmosterol acetates by thin-layer and column chromatography on silica gel G-silver nitrate, 72
 — Enrica G. Paoletti. Short communication re quantitative separation of C-27 sterol precursors of cholesterol, 84
 Gauslaa, Gerd. See Alec D. Keith
 Giovanelli, J. See A. R. Johnson
 Goswami, A. See J. G. Coniglio
 Gunstone, F. D. See W. W. Christie
 — See Howard W. Sprecher

H

- Hagemann, J. M., F. R. Earle, I. A. Wolff and A. S. Barclay. Search for new industrial oils. XIV. Seed oils of Labiatae, 371
 Hall, Michael O. See Mona E. Fewster
 Hamilton, James G. See Ronald C. Reitz
 Hansen, R. P. See R. G. Ackman
 Harlow, R. D. See Carter Litchfield
 Harris, R. A., E. R. Rivera, C. L. Villemez, Jr., and F. W. Quackenbush. Mechanism of suckling-rat hypercholesterolemia. II. Cholesterol biosynthesis and cholic acid turnover studies, 137
 Hatch, F. T., N. K. Freeman, L. C. Jensen, G. R. Stevens and F. T. Lindgren. Ultracentrifugal isolation of serum chylomicron-containing fractions with quantitation by infrared spectrometry and NCH elemental analysis, 183
 Heimberg, Murray. See Donald R. Van Harken
 Herodek, Sándor. The distribution of labeled palmitic acid into the diglycerides and triglycerides of rat adipose tissue, 299
 Hirota, K. See J. Van Der Veen
 Hoffmann, R. L. See C. D. Evans
 Holman, R. T. See Howard W. Sprecher
 Hopkins, C. Y., and R. Swingle. Eicosenoic acid and other fatty acids of *Sapindaceae* seed oils, 258
 Hopkins, S. M. See J. Tinoco
 Horrocks, L. A., and G. B. Ansell. The incorporation of ethanolamine into ether-containing lipids in rat brain, 329
 Horwitt, M. K. See L. A. Witting
 Howard, Charles F., Jr., and George W. Kittinger. Short communication re extraction of chromatographically isolated fatty acids from liquid scintillation fluid, 438
 Huang, T. C., and Arnis Kuksis. A comparative study of the lipids of globule membrane and fat core and of the milk serum of cows, 453
 — and Arnis Kuksis. A comparative study of the lipids of chylomicron membrane and fat core and of the lymph serum of dogs, 443
 Hunter, J. J. See D. A. Lutz
 Huttsell, T. C., and F. W. Quackenbush. Reduction of atherogenicity of natural fats by small additions of ethyl linoleate in the diet of the rat, 342

J

- Jangaard, P. M. See R. G. Ackman
 Jensen, L. C. See F. T. Hatch
 Jensen, R. G. See J. Sampugna
 Johnson, A. R. See E. Allen
 — K. E. Murray, A. C. Fogerty, B. H. Kennett, Judith A. Pearson and F. S. Shenstone. The reaction of methyl sterculate and malvalate with silver nitrate-silica gel and its use as a basis for the estimation of cyclopropane fatty acids, 316

- Judith A. Pearson, F. S. Shenstone, A. C. Fogerty and J. Giovanelli. The biosynthesis of cyclopropane and cyclopropene fatty acids in plant tissue, 308
 Johnson, P. See C. H. Burren
 Johnston, John M., G. Ananda Rao, Patricia A. Lowe and Barry E. Schwartz. The nature of the stimulatory role of the supernatant fraction of triglyceride synthesis by the α -glycerophosphate pathway, 14

K

- Kabara, Jon J., and N. T. Wertheissen. Brain cholesterol. XII. The incorporation of 1-¹⁴C-acetate into baboon sterol, 494
 Keith, Alec D., Gerd Gauslaa and Beverly S. Anderson. In vitro biosynthesis of fatty acids in *Drosophila melanogaster*, 429
 Kennett, B. H. See A. R. Johnson
 Kittinger, George W. See Charles F. Howard, Jr.
 Kleiman, R., V. L. Davison, F. R. Earle and H. J. Dutton. Determination of petroselinic acid by microreactor chromatography, 339
 — R. W. Miller, F. R. Earle and I. A. Wolff. (S)-1,2-Diacetyl-3-acetins: optically active triglycerides from *Euonymus verrucosus* seed oil, 473
 Kraml, M., J. Dubuc and D. Dvornik. Agents affecting lipid metabolism. XXVI. Specificity of some inhibitors of the late stages of cholesterol biosynthesis, 5
 Kritchevsky, Gene. See A. J. Bauman
 — See George Rouser
 Kuchmak, Myron. Short communication re contamination of methyl ester preparations during methylation of fatty acids on an ion exchange resin, 192
 Kuksis, Arnis, and L. Marai. Determination of the complete structure of natural lecithins, 217
 — See T. C. Huang
 Kuta, Edwin J., and Mary Yu. Polarography of conjugated unsaturated lipids, 411

L

- Law, John H. See James A. McCloskey
 Lepage, Marius. Identification and composition of turnip root lipids, 244
 Leveille, Gilbert A., Ronald S. Pardini and Jerry Ann Tillotson. Influence of medium-chain triglycerides on lipid metabolism in the chick, 461
 — Ronald S. Pardini and Jerry Ann Tillotson. Influence of medium-chain triglycerides on lipid metabolism in the rat, 287
 Likhite, V. N. See L. A. Witting
 Lindgren, F. T. See F. T. Hatch
 List, G. R. See C. D. Evans
 Litchfield, Carter, R. D. Harlow and Raymond Reiser. Gas-liquid chromatography of triglyceride mixtures containing both odd and even carbon number fatty acids, 363
 — Earlene Miller, R. D. Harlow and Raymond Reiser. The triglyceride composition of 17 seed fats rich in octanoic, decanoic or lauric acid, 345
 Lowe, Patricia A. See John M. Johnston
 Luthi, L. See S. Rous
 Lutz, D. A., C. R. Eddy and J. J. Hunter. X-ray diffraction study of some normal alkyl esters of long-chain acids, 204
 Lyman, R. L. See J. Tinoco

Mc

- McCloskey, James A., and John H. Law. Ring location in cyclopropane fatty acid esters by a mass spectrometric method, 225
 McConnell, D. G. See C. D. Evans
 McIntosh, D. J. See J. Tinoco

M

- Marai, L. See A. Kuksis
 Mead, James F. See Dorothy L. Fillerup
 Melin, Karl-Axel. See Lars-Ake Appleqvist
 Menzel, Daniel B. Short communication re reaction of oxidizing lipids with ribonuclease, 83
 Mercari, O., R. D. Peluffo and R. R. Brenner. Short communication re effect of insulin on the oxidative desaturation of α -linolenic, oleic and palmitic acids, 284

- Merdinger, Emanuel, and Charles E. Cwiakala. Short communication re fatty acids of the lipids from *Pullularia pullulans*, 276
- Mikolajczak, K. L., and C. R. Smith, Jr. Optically active trihydroxy acids of *Chamaepeuce* seed oils, 261
- and C. R. Smith, Jr. Pentacyclic triterpenes of *Jurinea analolica* Boiss. and *Jurinea consanguinea* DC, fruit 127
- Miller, Earline. See Carter Litchfield
- Miller, R. W. See R. Kleiman
- Mitchell, W. D., and M. J. Diver. Analysis of fecal neutral steroids and bile acids in humans on constant fat diet, 467
- Moore, J. L., T. Richardson and H. F. Deluca. Lysosomes and essential fatty acid deficiency, 8
- Morrison, W. R., and L. M. Smith. Fatty acid composition of milk phospholipids. II. Sheep, Indian buffalo and human milks, 178
- Mounts, T. L. See Ami Dolev
- Murray, K. E. See A. R. Johnson

N

- Narayan, K. Ananth. Short communication re disc electrophoresis of human and animal serum lipoproteins, 282
- Nawar, Wassef W. See Ann C. Noble
- Nelson, Gary J. The phospholipid composition of plasma in various mammalian species, 323
- Studies on the lipids of sheep red blood cells. I. Lipid composition in low and high potassium red cells, 64
- See George Rouser
- Nicolaides, N. The moncone and other wax alcohols of human skin surface lipid and their relation to the fatty acids of this lipid, 266
- Noble, Ann C., Charles Buziassy and Wassef W. Nawar. Short communication re thermal hydrolysis of some natural fats, 435
- Nutter, L. J. See O. S. Privett

O

- Olcott, H. S. See J. Van Der Veen
- O'Neill, John J. See Thomas E. Duffy
- Ottolenghi, Athos. Phospholipase activity of rat tissues and its modification by trypsin, 303

P

- Paoletti, Enrica G. See Giovanni Galli
- Pardini, Ronald S. See Gilbert A. Leveille
- Paulose, M. M. See S. Venkob Rao
- Pearson, Judith A. See E. Allen
- See A. R. Johnson
- Peluffo, R. D. See O. Mercuri
- Pereira, R. L. See C. Jane Wyatt
- Phillips, F., and C. V. Viswanathan. Short communication re a simple method of preparing alcohol acetates, wax esters and cholesterol esters, 437
- Piantadosi, Claude. See Shu-Sing Cheng
- Pitas, R. E. See J. Sampugna
- Powell, R. G., C. R. Smith, Jr., and I. A. Wolff. *cis-5, cis-9, cis-12*-Octadecatrienoic and some unusual oxygenated acids in *Xeranthemum annuum* seed oil, 172
- Privett, O. S., and L. J. Nutter. Determination of the structure of lecithins via the formation of acetylated 1,2 diglycerides, 149

Q

- Quackenbush, F. W. See R. A. Harris
- See T. C. Hutsell
- Quinn, J. G. See J. Sampugna

R

- Raju, P. K., and Raymond Reiser. Short communication re elimination of contaminants during the isolation of fatty acid esters by preparative gas-liquid chromatography, 197

- Rao, G. Ananda. See John M. Johnston
- Rao, S. Venkob, M. M. Paulose and B. Vijayalakshmi. Letter to the editor re a note on the fatty acids present in oilseed phospholipids, 88
- Re Cerconi, Daniela. See Claudio Galli
- Reiser, Raymond. See Robert E. Anderson
- See Nestor R. Bottino
- See Carter Litchfield
- See P. K. Raju
- Reitz, Ronald C., James G. Hamilton and Francis E. Cole. Fatty acid concentrations in synchronous cultures of *Chlorella pyrenoidosa*, grown in the presence and absence of glucose, 381
- Rhee, Ki Seon, R. R. Del Rosario and L. R. Dugan, Jr. Determination of plasmalogens after treating with a 2,4-dinitrophenylhydrazine-phosphoric acid reagent, 334
- Richardson, T. See J. L. Moore
- Rivera, E. R. See R. A. Harris
- Robertson, Alex, and Howard Sprecher. Human placental lipid metabolism. III. Synthesis and hydrolysis of phospholipids, 403
- Robinson, Ranga. See Eric Baer
- Rohwedder, W. K. See Ami Dolev
- Rous, S., L. Luthi and P. Favarger. Role of ATP in the inhibition of lipogenesis in fasted animals, 60
- Rouser, George. See A. J. Bauman
- See Manik L. Das
- See Gerald Simon
- Gene Kritchevsky, Gerald Simon and Gary J. Nelson. Quantitative analysis of brain and spinach leaf lipids employing allieic acid column chromatography and acetone for elution of glycolipids, 37

S

- Sampugna, J., J. G. Quinn, R. E. Pitas, D. L. Carpenter and R. G. Jensen. Digestion of butyrate glycerides by pancreatic lipase, 397
- Schulman, Jack H. See Dinesh O. Shah
- Schwartz, Barry E. See John M. Johnston
- Shah, Dinesh O., and Jack H. Schulman. Interaction of calcium ions with lecithin and sphingomyelin monolayers, 21
- Sheehan, G. See J. Tinoco
- Shenstone, F. S. See E. Allen
- See A. R. Johnson
- Siakotos, A. N. Short communication re rapid determination of lipids containing free amino groups with trinitrobenzene sulfonic acid reagent, 87
- See Thomas E. Duffy
- Simon, Gerald. See George Rouser
- and George Rouser. Phospholipids of the sea anemone: Quantitative distribution; absence of carbon-phosphorus linkages in glycerol phospholipids; structural elucidation of ceramide aminoethylphosphonate, 55
- Sipes, J. C. See R. G. Ackman
- Smith, C. R., Jr. See K. L. Mikolajczak
- See R. G. Powell
- Smith, L. M. See W. R. Morrison
- Snyder, Fred. See Shu-Sing Cheng
- See Randall Wood
- Sprecher, Howard W. See Alex Robertson
- H. J. Dutton, F. D. Gunstone, P. T. Sykes and R. T. Holman. Metabolic conversions and the positional distributions in liver lecithin of some unnatural dienoic acids in the rat, 122
- Stevens, G. R. See F. T. Hatch
- Storry, J. E. and B. Tuckley. Short communication re thin-layer chromatography of plasma lipids by single development, 501
- Swingle, R. See C. Y. Hopkins
- Sykes, P. T. See Howard W. Sprecher

T

- Theron, J. J. See L. A. Witting
- Tillotson, Jerry Ann. See Gilbert A. Leveille
- Tinoco, J., S. M. Hopkins, D. J. McIntosh, G. Sheehan and R. L. Lyman. Fractionation and analysis of rat liver ^{14}C -lecithins labeled in vivo, 479
- Tsao, Shuang-Shine, and W. E. Cornatzer. Biosynthesis of phospholipids in subcellular particles from cultured cells of human tissue, 424
- and W. E. Cornatzer. Chemical composition of subcellular particles from cultured cells of human tissue, 41
- Tuckley, B. See J. E. Storry

V

- Van Deenen, L. L. M. See H. Van Den Bosch
 Vandenberg, Gloria A. See Nestor R. Bottino
 Van Den Bosch, H., H. M. Van Der Elzen and L. L. M. Van Deenen. Short communication re the phospholipases of yeast, 279
 Van Der Elzen, H. M. See H. Van Den Bosch
 Van Der Veen, J., K. Hirota and H. S. Olcott. Comparison of column chromatographic methods for the quantitative determination of mono- and diagalactodiglycerides in fresh alfalfa (*Medicago sativa*), 406
 Van Harken, Donald R., Thomas O. Brown and Murray Heimberg. Hepatic lipid metabolism in experimental diabetes: III. Synthesis and utilization of triglycerides, 231
 Vijayalakshmi, B. See S. Venkob Rao
 Villemex, C. L., Jr. See R. A. Harris
 Viswanathan, C. V. See F. Phillips

W

- Walker, Brian L. Maternal diet and brain fatty acids in young rats, 497
 Werthessen, N. T. See Jon J. Kabara
 Wishner, Lawrence A. See Nancy M. Derrick
 Witting, L. A. The effect of antioxidant deficiency on tissue lipid composition in the rat. IV. Peroxidation and interconversion of polyunsaturated fatty acids in mus-

- cle phospholipids, 109
 — and M. K. Horwitt. The effect of antioxidant deficiency on tissue lipid composition in the rat. I. Gastrocnemius and quadriceps muscle, 89
 — V. N. Likhite and M. K. Horwitt. The effect of antioxidant deficiency on tissue lipid composition in the rat. III. Testes, 103
 — J. J. Theron and M. K. Horwitt. The effect of antioxidant deficiency on tissue lipid composition in the rat. II. Liver, 97
 Wolff, I. A. See J. M. Hagemann
 — See R. Kleiman
 — See R. G. Powell
 Wood, Randall. GLC and TLC analysis of isopropylidene derivatives of isomeric polyhydroxy acids derived from positional and geometrical isomers of unsaturated fatty acids, 199
 — and Fred Snyder. Chemical and physical properties of isomeric glyceryl monoethers, 161
 Wright, Lemuel D. Short communication re an in vivo approach to the mechanism of cholesterol deposition in vivo, 277
 Wyatt, C. Jane, R. L. Pereira and E. A. Day. The lactone precursor in fresh milk fat: Identification of the hydroxy fatty acids, 208

X-Y-Z

- Yu, Mary. See Edwin J. Kuta

SUBJECT

A

- Acanthosyrus spinescens**, triglyceride profile, 363
- Acetic acid**, acetates of pentacyclic triterpene alcohols, 127
- incorporation into baboon brain cholesterol, 494
 - incorporation into fatty acids and cholesterol in chicken liver, effect of diet, 461
 - incorporation into unsaturated fatty acids in antioxidant deficiency, 89, 97
 - oleyl alcohol ester, preparative procedure, melting point, 437
 - as precursor for fatty acids in *Drosophila melanogaster* larvae, 429
- Acetolysis**, of phosphatidyl choline, acyl migrations, 149
- Acetylation**, of diglycerides, 217
- of sterols, 72
 - of triterpene alcohols, 127
- Acetyl-Coenzyme A**, failure to inhibit acylation in intestinal mucosa, 14
- Acetyl 1,2-diacylglycerol**, structure dtmn, 149, 217
- ACP**. See acyl carrier protein
- Actinodaphne hookeri**, fatty acid profile; triglyceride profile, 345
- Actinolema macrolema Bioss.**, petroselinic acid content, 339
- Acyl carrier protein-ACP**, implicated in triglyceride biosynthesis, 14
- Acyl-coenzyme A**: acyl-glycerolphosphorylethanolamine acyltransferase-EC 2.3.1., in human placenta, 403
- Acyl migration**, during acetolysis of phosphatidyl choline, 149
- 1-(O-phosphoethanolamine)-N-Acyl sphingosine**. See ceramide aminoethylphosphonate
- Acyltransferase-EC-2.3.1.** — acyl Co A: acyl GPE transferase, human placenta, 403
- in human placenta, 403
- Adenosine triphosphate**, role in lipogenesis, 60
- Adipose tissue**, fatty acid profile, 212
- lipids in vitamin E deficient rats, 133
 - rat, fatty acid synthesis and diet, 287
 - rat, palmitic acid incorporation into di- and triglycerides, 299
- Adonis aestivalis L.**, diacyl-3-acetin content, 473
- Adrenals**, baboon, acetic acid incorporation into cholesterol, 494
- Adsorption isotherms**, for cholesterol on Permutit, 277
- Agastache urticifolia (Benth.) O. Ktze.**, fatty acid profile, 371
- Age**, effect on human brain lipids, 295
- Ajuga chia (Poir.) Schreb.**, fatty acid profile, 371
- Ajuga iva Schreb.**, fatty acid profile, 371
- Akebia quinata Decne.**, diacyl-3-acetin content, 473
- Akebia trifoliata (Thunb.) Koidz.**, diacyl-3-acetin content, 473
- Albumin**, as carrier of lipids in electrophoresis, 282
- Alcohol**, pentacyclic triterpenes, 127
- Alcohol acetates**, preparative procedure, 437
- Alcohol-aliphatic**, iso- and anteiso, 266
- chain elongation, 266
- Alcohol-higher aliphatic**, in human skin surface lipids, 266
- nomenclature, short-hand, 266
 - polyenes, in human skin surface lipids, 266
 - position isomers, of human skin surface lipid, 266
- Alfalfa (Medicago sativa)**, galactoglycerides, 406
- Alkyl acyl glyceryl-3-phosphorylcholine**, ethanolamine incorporation in rat brain, 329
- Alkyl acyl glyceryl-3-phosphorylethanolamine**, ethanolamine incorporation in rat brain, 329
- Alloxan**, experimental diabetics in rats, 231
- Alloxan-diabetic rat**, oxidative desaturation of fatty acids, 284
- Aldehydes**, alkanals and alk-2-enals accumulation in vitamin E deficient rats, 133
- unsaturated, borohydride titration, 280
 - unsaturated, hydrogen iodine values, 280
- Amethystea caerulea L.**, fatty acid profile, 371
- Amine containing lipids**, analysis by trinitrobenzene sulfonic acid, 87
- 2-Aminoethylphosphonic acid**, in sea anemone phosphonolipids, 55
- 20 α -(2-dimethylaminoethyl) Amino-5 α -pregnan-3 β -ol dihydrochloride**, 22, 25-DAC (SC-11952), cholesterol biosynthesis inhibitor, 5
- α -Amyrin**, in *Jurinea* fruit, 127

- melting point, physical properties, 127
- β -Amyrin**, in *Jurinea* fruit, 127
- melting point, physical properties, 127
- β -Amyrin acetate**, melting point, physical properties, 127
- Anisomeles indica (L.) O. Ktze**, fatty acid profile, 371
- Anthopleura elegantissima**, sea anemone, phospholipids, 55
- Antibiotics**, in rat diet, 155
- Antioxidant deficiency**, effect on rat liver phospholipid fatty acid comp, 97
- effect on rat muscle phospholipid fatty acid comp, 89, 109
 - effect on rat testes phospholipid fatty acid comp, 103
 - fluorescent pigment accumulations in rat liver, 97
 - fluorescent pigment accumulation in rat muscle, 89
 - incorporation of acetic acid into rat lipids, 89, 97, 103
 - rat, effect of dietary fat comp, 89
 - rat, relieved by N,N'-diphenyl-p-phenylenediamine, 89, 97
 - rat, relieved by sodium selenide and methionine, 89
 - subcellular alterations in rats, 97
 - See also α -tocopherol deficiency
- Arachidic acid**, 258
- in castor seed phospholipids, 88
- Arachidonic acid**, in castor seed phospholipids, 88
- decrease in alloxan-diabetic rats, 284
 - incorporation into rat tissues, 114
 - increase in rat liver phospholipids in antioxidant deficiency, 97
 - increase in rat testes phospholipids in antioxidant deficiency, 103
 - metabolism in rat, pyridoxine deficiency, 484
 - methyl ester, in rat diet, 155
 - in phosphatidyl choline, 149
 - in rat muscle phospholipids, increase during antioxidant deficiency, 89, 109
 - synthesized from linoleic acid, 114
 - in Triton hyperlipemia, 212
- Arachidonic acid (20:4 ω 6)**, GLC, 502
- in phosphatidyl choline of rat liver, 122
- Atherogenesis**, *in vitro* model, 277
- Atherosclerosis**, *in vitro* model, 277
- rat, inhibited by dietary ethyl linoleate, 342
- Automatic sample injector**, for GLC, 351
- Autoxidation**, of lipids, prevention, 1
- of methyl linoleate, 28
 - of methyl linoleate, failure of thiobarbituric acid test, 133
 - of tissue lipids in vitamin E deficient rats, 133
- AY-9944**. See *trans*-1,4-bis-(2-chlorobenzylaminoethyl) cyclohexane dihydrochloride
- Azelaldehydic acid**, methyl ester, hydrogen iodine values, 280

B

- Baboon adrenals**, acetic acid incorporation into cholesterol, 494
- Baboon brain**, acetic acid incorporation into cholesterol, 494
- Baboon kidney**, acetic acid incorporation into cholesterol, 494
- Baboon liver**, acetic acid incorporation into cholesterol, 494
- Baboon spleen**, acetic acid incorporation into cholesterol, 494
- Ballota acetabulosa Benth.**, fatty acid profile, 371
- Ballota hispanica (L.) Benth.**, fatty acid profile, 371
- Ballota nigra L. ssp. ruderalis (Sw.) Briq.**, fatty acid profile, 371
- Batyl alcohol**, GLC, 143
- from tumor lipids, 143
- Betonica alopecurus L.**, fatty acid profile, 371
- Bile acids**, control of cholesterol biosynthesis through a feedback mechanism, 137
- dietary inhibit cholesterol biosynthesis, 137
 - excretion by humans, 467
- Bile acid infusion**, effect on cholesterol biosynthesis in suckling rats, 137
- Bile duct ligation**, effect on cholesterol biosynthesis, 137
- Blood plasma**, bovine, TLC, 501
- phospholipids of various mammals, 323
- Borohydride**, to titrate double bonds, 280
- Bovine milk**, comparison of summer and winter, 443
- lipids of serum and membranes, 443
 - protein and lipid relationships, 443

- Bovine blood, plasma lipids, TLC, 501
 Bovine blood plasma, phospholipid comp, 323
 Bovine heart ventricle, lipids comp, 1
 Brachychiton acerifolium, fatty acid, cyclopropane and cyclopropene, profile of seeds, 308
 — fatty acid profile of seed lipids, 316
 — seed lipids, profile and biosynthesis, 308
 Brachychiton populneum, fatty acid, cyclopropane and cyclopropene, profile of seeds, 308
 — fatty acid profile of seed lipids, 316
 — seed lipids, profile and biosynthesis, 308
 Brain, baboon, acetic acid incorporation into cholesterol, 494
 — glycolipids, 37
 Brain, rat, development, 76
 — ethanolamine incorporation into lipid ethers, 329
 — incorporation of 1-¹⁴C-arachidonic acid radio activity, 484
 Brain sterols, during growth of rats, 76
 Brain weight, and total lipids during growth of rats, 76
 Brasoria scutellarioides Engelm. and Gray., fatty acid profile, 371
 Brevoortia tyrannus (menhaden), triglyceride profile, 363
 Eupleurum croceum Fenzl., petroselinic acid content, 339
 Butanal, in vitamin E deficient rats, 133
 Butyryl-1,3-dipalmitin, hydrolysis by pancreatic lipase, 397
 — synthesis, 397
 Butyryl-2,3-dipalmitin, hydrolysis by pancreatic lipase, 397
 — synthesis, 397
 1-Butyryl-3-palmitin, hydrolysis by pancreatic lipase, 397
 — synthesis, 397

C

- Calamintha clinopodium Benth., fatty acid profile, 371
 Calamintha nepetoides Jord., fatty acid profile, 371
 Calcium, interaction with phosphatidyl choline and sphingomyelin monolayers, 21
 Campesterol glucoside, in turnip root, 244
 Carbonyl analysis, method, 133
 Carbonyl compounds, isolated from tissue lipids, 133
 Carcinoma, Ehrlich ascites cells, glycerol ether diesters, 193
 — human cervix lipids in tissue cultured HeLa cells, 41
 — human nasopharynx, lipids in tissue cultured KB cells, 41
 Cardiolipid, in turnip roots, 244
 Cardiospermum canescens Wall., fatty acid profile, 258
 Cardiospermum grandiflorum Sw., fatty acid profile, 258
 Cardiospermum haliacabum, fatty acid profile, 258
 β -Carotene, polarography, 411
 — in turnip roots, 244
 γ -Carotene, in turnip roots, 244
 Carotenoids, in Pullularia pullulana, 276
 Carotenoids, in turnip roots, 244
 Castor seed, fatty acid profile of phospholipids, 88
 Cat blood plasma, phospholipid comp, 323
 Cathepsin, subcellular distribution in essential fatty acid deficiency in rats, 8
 Celastrus orbiculatus Thunb., diacyl-3-acetin content, 473
 Celastrus scandens L., diacyl-3-acetin content, 473
 Cerebral cortex, dog, lipid synthesis *in vitro*, 195
 Cerebrosides, during rat brain development, 76
 — human brain, 37
 — in human brain, age, 295
 — isolation procedure, 37
 Cerebroside sulfate, in human brain, age, 295
 Ceramide, in human brain, age, 295
 Ceramide aminoethylphosphonate, fatty acid profiles, sen anemone, 55
 — hydrolysis, 55
 — oxidative degradation, 55
 — in sea anemone, 55
 — structural dtmn, 55
 Chamaepeuce afra (Jacq.) DC., 261
 Chamaepeuce hispanica DC., 261
 Chicken, influence of triglycerides (8:0 and 10:0) on lipid metabolism, 461
 Chicken liver, fatty acid Δ 9,10-desaturase activity, 419
 Chicken serum, lipoproteins by disc electrophoresis, 282
 Chimyl alcohol, GLC, 143
 — from tumor lipids, 143
 Chlorella pyrenoidosa, fatty acid profile and lipid extraction of, 381
 — See microorganisms

LIPIDS, VOL. 2, No. 6

- Δ 5,7-Cholestadiene-3 β -ol. See 7-Dehydrocholesterol
 Δ 5,24-Cholestadiene-3 β -ol. See Desmosterol
 Δ 5,7,24-Cholestatriene-3 β -ol. See 7-Dehydrodesmosterol
 Δ 5-Cholesten-3 β -ol. See Cholesterol
 Cholesterol, acetic acid incorporation into baboon brain, 494
 — adsorption on Permutit, 277
 — biosynthesis, feedback control by bile acids, 137
 — biosynthesis, in suckling rat, 137
 — bovine heart ventricle, 1
 — in bovine milk fat globules, 443
 — in bovine milk membranes, 443
 — in bovine milk serum, 443
 — in chicken liver, effect of diet, 461
 — DEAE column chromatography, 1
 — in dog lymph chylomicron membranes, 453
 — in dog lymph fat cores, 453
 — in dog lymph serum, 453
 — dtmn in human feces, 467
 — excreted by *Fasciola hepatica* L., 353
 — in human brain, age, 295
 — in human skin surface lipids, 266
 — incorporation of glucose carbon in dog brain, 195
 — infrared spectrum, 64
 — phospholipid ratio in erythrocytes, 64
 — separation from desmosterol, 72
 — separation from C-27 sterol precursors, 84
 — sheep erythrocytes, 64
 — TLC, 501
 — in Triton hyperlipemia, 212
 Cholesterol acetate, in human skin surface lipids, 266
 Cholesterol biosynthesis, inhibitors of, 5
 Cholesterol esters, in dog lymph serum, 453
 — excreted by *Fasciola hepatica* L., 353
 — incorporation of glucose carbon in dog brain, 195
 — preparative procedure, 437
 — rat liver, incorporation of 1-¹⁴C-arachidonic acid radio activity, 484
 — TLC, 501
 — in Triton hyperlipemia, 212
 Cholesterol levels, in rat liver and blood serum, effect of dietary triglycerides, 287
 Cholesteryl palmitoleate, preparative procedure, melting point, 437
 Cholic acid, excretion in suckling rats, 137
 — turnover in suckling rat, 137
 Choline, microbiological assay, 239
 Choline deficiency, in rats, 239
 Chromatography, column, aliphatic alcohol acetates, 266
 — alumina, α -tocopherol oxidation products, 47
 — alumina for α -tocopherol and α -tocopherol quinone, 390
 — Amberlite as source of contamination of methyl esters, 192
 — carotenoids, 244
 — Celite-Silica Gel G impregnated with silver nitrate, column prep, 72
 — Celite-Silica Gel G impregnated with silver nitrate for sterol acetates, 72, 84
 — charcoal-Celite (Nuchar-Attaclay) for galactoglycerides, 406
 — DEAE of bovine heart ventricle, 1
 — DEAE of human brain, 295
 — 2,4-dinitrophenylhydrazine column to determine carbonyl compounds, 133
 — Dowex for analysis of neutral steroids and bile acids, 467
 — gradient elution of phospholipids, 64
 — methyl 9-hydroxystearate, 28
 — methyl 13-hydroxystearate, 28
 — methyl ketostearate, 28
 — methyl stearate, 28
 — phospholipids, 64
 — separation of phospholipids from triglycerides, 114
 — silicic acid for diacyl-3-acetins, 473
 — silicic acid for galactoglycerides, 406
 — silicic acid for glycolipids, 37
 — solvents for sterol acetates, 72
 Chromatography, gas-liquid, acetyl diacylglycerol, 217
 — analytical conditions for fatty acid esters, 502
 — analytical conditions for triglycerides, 363
 — of batyl alcohol, 143
 — of chimyl alcohol, 143
 — column packing (unsaturated fatty acid esters), 502
 — correction factors for quantitation of polyunsaturated fatty acids, 251
 — *Euonymus verrucosus* seed oil, 473
 — fatty acid analysis, 479
 — fatty acid methyl ester, 217
 — of fish oils, triglycerides, 363
 — glycerol ethers, mono-, analytical conditions, 161

- hydroxy fatty acid chain length related to retention time, 206
- hydroxy fatty acid methyl ester-TMS, retention times, 208
- isopropylidenes from isomeric polyhydroxy fatty acids, 199
- methyl hydroxystearates, 28
- open tubular for polyunsaturated fatty acids, quantitation problem, 251
- of pentacyclic triterpene acetates from *Jurinea* fruit, 127
- phytanic acid diastereoisomers, 357
- preparative for acetyl diacylglycerol, 217
- preparative, for fatty acid esters, 197
- preparative, monoene alcohol acetates from skin surface lipids, 266
- pristanic acid diastereoisomers, 357
- sample injection system, automatic solvent-free, 351
- of seed oils, triglycerides, 363
- triglycerides, 363, 473
- of tumor lipids, 143
- of unsaturated fatty acid esters, 502
- Chromatography, gas-liquid-mass spectrometer, dtmn of branching in acid esters, 225**
- dtmn of cyclopropane fatty acids, 225
- Chromatography, mass spectrometer. See Chromatography, gas-liquid-mass spectrometer**
- Chromatography, paper, paraffin-coated, α -tocopherol oxidation products, 47**
 - silicic acid-impregnated glass paper (phospholipids), 41
 - solvent system (dinitrophenyl hydrazones), 484
 - solvent system for glass fiber paper (fatty acids), 381
 - solvent system for paraffin impregnated (α -tocopherol and derivatives), 390
 - solvent system for silicic acid impregnated (α -tocopherol and derivatives), 390
- Chromatography, Sephadex, of lipids, 323**
 - removal of non-lipid contaminants, 37
 - G-50 to remove CoA, 14
 - of ribonuclease, 88
- Chromatography, thin-layer, blood plasma lipids, 501**
 - cyclopropane fatty acids on silica gel-silver nitrate, 308, 316
 - cyclopropane fatty acids on silica gel-silver nitrate, 308, 316
 - detection procedures, 467
 - developing chamber, 501
 - glycolipids, 37
 - isopropylidenes from isomeric polyhydroxy fatty acids, 199
 - pentacyclic triterpene acetates, 127
 - phospholipids, 37, 64
 - polyhydroxy stearic acids, 261
 - polyunsaturated fatty acid methyl esters by chain length and extent and location of unsaturation, 89
 - plate prep (silica gel), 501
 - plate prep (silica gel-silver nitrate), 339, 479
 - preparative (silica-gel-silver nitrate), 127
 - preparative (triglycerides), 345, 489
 - preparative (phosphatidyl choline), 479
 - preparative, lysophosphatidyl choline and fatty acids, 122
 - quantitative, 1, 37
 - Rf values for α -tocopherol and derivatives, 390
 - Saran Wrap, protection from iodine vapors, 244
 - separation of phospholipids from triglycerides, 114
 - similarity in migration of phthalic acid esters and triglycerides, 85
 - solvent system (acetyl diacylglycerol), 217
 - solvent system (bile acids), 467
 - solvent system (carotenoids), 244
 - solvent system (cyclopropane and cyclopropane fatty acids), 316
 - solvent system (diacyl-3-acetin), 473
 - solvent system (diacylglycerol ethers), 143
 - solvent system (glyceride), 299
 - solvent system (glycerol ethers), 161
 - solvent system (glycolipids and phospholipids), 295
 - solvent system (methyl petroselinates), 339
 - solvent system (neutral lipids), 244, 295, 353, 453
 - solvent system (neutral lipids, phospholipids), 212
 - solvent system (neutral and polar lipids), 14
 - solvent system (pentacyclic triterpene acetates), 127
 - solvent system (phosphatidyl choline), 479
 - solvent system (phospholipids), 323, 453
 - solvent system (phospholipids, glycerides, cholesterol esters), 501
 - solvent system (polar lipids), 244, 353
 - solvent system (polyhydroxy fatty acids), 261
 - solvent system (polyunsaturated fatty acids), 89
 - solvent system (α -tocopherol, α -tocopherol quinone, α -tocopheronic acid), 390
 - solvent system (triglycerides), 149
 - sterol acetates, 72
 - α -tocopherol oxidation products, 47
 - triglycerides by numbers of double bonds, 149
 - two-dimensional, glycolipids, 37
 - two-dimensional, phospholipids, 37, 64, 323
 - two-dimensional, phospholipids, phosphonolipids, 55
- Chylomicron, concentration in serum, 183**
 - dog, isolation from lymph, 453
 - isolation from serum by ultracentrifugation, 183
 - membrane prep from bovine milk, 443
 - membrane prep from dog lymph serum, 453
 - protein content, 183
 - quantitation by elemental analysis, 183
 - transport of linoleic acid, 114
- Cinnamomum camphora, fatty acid profile; triglyceride profile, 345**
- Cleonia lusitana L., fatty acid profile, 371**
- Clostridium perfringens, phospholipase C (Type A toxin), 149**
- CoA. See coenzyme A**
- Coconut oil, fatty acid profile of, 287**
 - solvent for cholesterol-Permutit adsorption isotherm, 277
- Cod, *Gadus morhua*, liver oil fatty acid profile, 251**
- Coenzyme A, in biosynthesis of triglycerides, 14**
- Compound O, α -tocopherol oxidation product, 47**
- Contaminants, prevention during preparative GLC, 197**
 - of lipid extracts, by plastic bag phthalic acid esters, 85
- Contamination, of methyl ester by ion exchange resins, 192**
- Coprostanol, dtmn in human feces, 467**
- Coprostanone, dtmn in human feces, 467**
- Corn oil, fatty acid profile of, 287**
 - solvent for cholesterol-Permutit adsorption isotherm, 277
- Coronarinic acid. See cis-9-L,10-L-epoxy-cis-12-octadecadecenoic acid**
- Correlation matrix, of lipid fractions and cholesterol ester fatty acids, 212**
 - of lipid fractions and phospholipid fatty acids, 212
 - of lipid fractions and triglyceride fatty acids, 212
- Counterstream distribution, solvent system for cis-9-L,10-L-epoxy-cis-12-octadecanoic acid methyl esters, 172**
 - solvent systems for cis-9,10-epoxyoctadecanoic acid methyl esters, 172
 - solvent systems for hydroxy acid esters, 172
 - solvent system for cis-5, cis-9, cis-12-octadecatrienoic acid methyl ester, 172
- Cream, raw sweet, polar glyceride, 208**
- Creatinuria, in rat antioxidant deficiency, 89, 109**
- Crotalus adamanteus, phospholipase A, 122**
- Crotonaldehyde, hydrogen iodine value, 280**
- Cruciferae, turnip roots, Laurentian and Wye, 244**
- Cryptexanthin, in turnip roots, 244**
- Cuphea carthagenensis, fatty acid profile; triglyceride profile, 345**
- Cuphea hookeriana, fatty acid profile; triglyceride profile, 345**
- Cuphea ignea, fatty acid profile; triglyceride profile, 345**
- Cuphea lanceolata, fatty acid profile; triglyceride profile, 345**
- Cuphea flava, fatty acid profile; triglyceride profile, 345**
- Cuphea painteri, fatty acid profile; triglyceride profile, 345**
- trans-1,4-bis(2-chlorobenzylaminomethyl)Cyclohexane dihydrochloride, AY-9944, cholesterol biosynthesis inhibitor, 5**
- trans-1,4-bis(2-dichlorobenzylaminomethyl)Cyclohexane dihydrochloride, AY 9944, cholesterol biosynthesis inhibitor, 84**
- Cyclopropane fatty acid, ring location, 225**

D

- Daucus guttatus Sibth. & Sm., petroselinic acid content, 339**
- n-Decyl alcohol, from plastic bag contaminants, 85**
- 7-Dehydrocholesterol, accumulation in liver by inhibitors, 84**
 - cholesterol biosynthesis, 5
- 24-Dehydrocholesterol. See desmosterol**
- 7-Dehydrocholesterol-37-reductase. See 37-reductase**

- 7-Dehydrodesmosterol**, accumulation in liver by inhibitors, 84
- 24-Dehydrosterol- Δ^{24} -reductase**. See Δ^{24} -reductase
- Deoxycholic acid**, dtm in human feces, 467
- Deoxyribonucleic acid**, in cultured human cells, 41
- Desmosterol**, cholesterol biosynthesis, 5
— separation from cholesterol, 72
— See also 24-dehydrocholesterol
- Development of rat brain**, lipid changes in rat brain development, 76
- Diabetic rats**, alloxan, oxidative desaturation of fatty acids, 284
- Diacyl-3-acetin**, spectrum, NMR, 473
(S)-1,2-Diacyl-3-acetin, in seed oils, 473
— spectrum, NMR, 473
— synthetic procedure, 473
- Diacylglycerol ethers**, in human lymphosarcoma, 143
— in Walker 256 carcinoma, 143
- (S)-1,2-Distearo-3-acetin, synthetic procedure, 473
- N, N'-Dibenzylethylenediamine**, cholesterol biosynthesis inhibitor, 5
- 5,6-Dibromocholestan- β -ol**, derivative of cholesterol, 5
- Dibutyryl**, hydrolysis by pancreatic lipase, 397
- Didecyl phthalate**, contaminate of plastic bags, 85
— infrared spectrum, 85, 198
- Diet**, chicken, coconut oil effect on lipid metabolism, 461
— chicken, corn oil effect on lipid metabolism, 461
— chicken, triglycerides (8:0 and 10:0) effect on lipid metabolism, 461
— cod-liver oil, vitamin E deficient, 133
— corn oil and coconut oil effect on cholesterol levels, 287
— effect of corn oil and butterfat on dog lymph lipids, 453
— effect of plant sterols on dog lymph lipids, 453
— fat-free, rats, 155
— fatty acid synthesis in adipose tissue of rat, 287
— high fat, in suckling rats hypercholesterolemia, 137
— maternal, effect on brain fatty acids in rats, 497
— production of atherosclerosis in rats, 342
— pyridoxine deficient, for rats, 484
— triglycerides (8:0 and 10:0) effect on cholesterol levels, 287
- Dietary fat**, fatty acid profiles, 342
- Digalactoglycerides**, in alfalfa, 406
— infrared spectrum, 406
- Diglycerides**, in bovine milk fat globules, 443
— in bovine milk membranes, 443
— in bovine milk serum, 443
— in dog lymph chylomicron membranes, 453
— in dog lymph fat cores, 453
— in dog lymph serum, 453
— fatty acid profile in bovine milk, 443
— fatty acid profile of dog lymph, 453
— palmitic acid incorporation by rat adipose tissue, 299
— prep by pancreatic lipase, 440
— structure dtm by acetylation, 149, 217
— 1,3-from 1,2-by TLC, 501
- 1,2-Diglyceride**, biosynthesis in intestinal mucosa, 14
- Diglycerophosphoryl glycerol**, from turnip roots, 244
- Diglycosyl diglyceride**, spinach leaf, 37
- Dihydromalvalic acid**, biosynthesis in plant tissues, 308
— See 8,9-methyleneheptadecanoic acid
- Dihydrostercolic acid**, biosynthesis in plant tissues, 308
— lack of effect on fatty acid $\Delta^9,10$ -desaturase, 419
— See 9,10-methyleneoctadecanoic acid
- Dinitrophenylhydrazones**, to determine plasmalogens, 334
- 1,2-Dioctadecen-3-eicosapentaeno-**, hydrolysis by pancreatic lipase, 489
- 1,2-Dioctadeceno-3-eicosapentaenin**, hydrolysis by pancreatic lipase, 489
- 1,2-Dioctadecenyl-3-eicosapentaenyl glycerol**, hydrolysis by pancreatic lipase, 489
- Diethyl phthalate**, contaminate of plastic bags, 85
- L- α -Dipalmitoyl glycerophosphoryl choline**, calcium interaction with monolayer, 21
— monolayer properties, 21
- N, N'-Diphenyl-p-phenylenediamine**, use as dietary antioxidant, 89, 97
- Diphosphatidyl glycerol**, bovine heart ventricle, 1
— DEAE column chromatography, 1
— See cardiolipin
- Dipoles**, of sphingomyelin, 21
- (S)-1,2-Distearo-3-acetin, spectrum, optical rotatory dispersion, 473
- Docosahexaenoic acid**, decrease in rat muscle phospholipids in antioxidant deficiency, 109
— decrease in alloxan-diabetic rats, 284
— in rat muscle phospholipids in antioxidant deficiency, 89
- Docosanoic acid** (20:0), GLC, 502
- Docosapentaenoic acid**, in rat muscle phospholipids in antioxidant deficiency, 89
- Docosenoic acid** (22:1), GLC, 502
- Docos-12-en-1-ol**, in human skin lipids, 266
- Dodones**, fatty acid profile, 258
- Dodones lanceolata** F. Meull., fatty acid profile, 258
- Dodones viscosa** Jacq., fatty acid profile, 258
- Dog**, lipid synthesis in brain, 195
— experimental endogenous hyperlipemia, 212
— lymph, lipids of serum and chylomicrons, 453
— lymph, protein and lipid relationships in chylomicron membranes, 453
- Dog blood plasma**, phospholipid comp, 323
- Double bond**, hydroxylation by permanganate, 199
- Double bond conjugation**, polarography, 411
- Dracocephalum moldavica** L., fatty acid profile, 371
- Drosophila melanogaster** larvae, fatty acid biosynthesis, 429
— growth and collection, 429
- Ducrosia anethifolia** Boiss., petroselinic acid content, 339

E

- EC 3.1.2.2**. See palmitoyl-coenzyme A hydrolase
- EC 3.1.3.4**. See L- α -phosphatidate phosphohydrolase
- EFA**. See fatty acid, unsaturated
- Egg**, phosphatidyl choline structure, 149
— phosphatidyl choline, reference in analysis of turnip root lipids, 244
- Egg yolk**, phosphatidyl choline, 217
— phosphatidyl choline in fatty acid profiles, 217
- Ehrlich ascites carcinoma cells**, glycerol ether diesters, 193
- 7,13-Eicosadienoic acid**, synthesis, metabolism in rat, 122
- 7,13-Eicosadiynoic acid**, methyl ester, reduction to methyl 7,13-eicosadienoate, 122
- Eicosanoic acid** (20:0), GLC, 502
- Eicosapentaenoic acid**, decrease in alloxan-diabetic rats, 284
— decrease in rat liver phospholipids in antioxidant deficiency, 97
- Eicosatetraenoic acid**, from linoleic acid, in rat intestine, 155
- Eicosatrienoic acid**, decrease in alloxan-diabetic rats, 284
— in rat mucosa lipids, 155
- Eicosa-5,8,11-trienoic acid**, in rat phospholipids, replacement by arachidonic acid, 114
- Eicosenoic acid**, in *Sapindaceae* seed oils, 258
- 11-Eicosenoic acid**, isopropylidenes of, GLC, TLC, 199
- cis-5-Eicosenoic acid**, in synthesis of cyclopropane fatty acids, 225
- cis-11-Eicosenoic acid**, from *Koeleruteria* and *Cardiospermum*, 258
- Eicos-10-en-1-ol**, in human skin lipids, 266
- Elaidic acid**, isopropylidenes of, GLC, TLC, 199
- Electron microscopy**, liver in antioxidant deficiency, 97
- Electrophoresis**, of aqueous dispersion of sphingomyelin, 21
— disc, of serum lipoproteins, 282
— disc, serum lipoprotein patterns of chicken, mouse, rat, rabbit and human, 282
- Eleostearic acid**, polarography, 411
— See 9,11,13-octadecatrienoic acid
- Elsholtzia patrinii** Garcke, fatty acid profile, 371
- Elsholtzia splendens** Nak., fatty acid profile, 371
- Encephalomalacia**, in the chick, resemblances to dietary antioxidant deficiency, 103
- Enterohepatic recirculation**, of bile, effect on cholesterol biosynthesis in suckling rat, 137
- Enthymus alletteratus** (tuna), triglyceride profile, 363
- Enzyme kinetic data**, K_m, L- α -glycerophosphate, 14
— K_m, 2-monopalmitin, 14
- Epididymal fat pad**, rat, fatty acid profile of triglycerides and phospholipids, 114
— in vitamin E deficient rats, 133
- Epoxy acids**, in *Xeranthemum annuum* seed oil, 172
- cis-9-L, 10-L-Epoxyoctadecenoic acid**, in *Xeranthemum annuum* seed oil, 172
- cis-9-L, 10-L-Epoxy-cis-12-octadecenoic acid** (coronarin), in *Xeranthemum annuum* seed oil, 172
- cis-12,13-Epoxy-cis-9-octadecenoic acid**, hydroxylation, nomenclature of isomers, 286
- cis-12-D, 13-D-Epoxy-cis-9-octadecenoic acid** (vernolic), in *Xeranthemum annuum* seed oil, 172
- Eremostachys laciniata** (L.) Bunge, fatty acid profile, 371
- Eremostachys speciosa**, Rupr., fatty acid profile, 371
- Eremostachys vicaryi** Benth., fatty acid profile, 371

- Ergosterol**, polarography, 411
- Eruic acid**, isopropylidenes of, GLC, TLC, 199
- in rape and mustard seed, 88
- Eryngium cheiranthium** Decne., petroselinic acid content, 339
- Erythrocyte**, isolation, 64
- sheep, lipid comp, 64
- Essential fatty acid deficiency**, in rats, 8
- Esters**, n-alkyl of fatty acids, 204
- homologous series related to x-ray long-spacings, 204
- Ethane**, from linoleic acid autooxidation, 432
- Ethanolamine**, analysis by trinitrobenzene sulfonic acid, 87
- incorporation into lipid ethers in rat brain, 329
- Ether**, glycerol, synthesis, 161
- glycerol lipids containing ethanolamine in rat brain, 329
- See also parent compound
- 1,2-propanediol 3-octadecyl, diacetate, infrared spectrum, 161
- 1, 2-propanediol 3-octadecyl, ditrifluoroacetate, infrared spectrum, 161
- 1,2-propanediol 3-octadecyl, racemic, infrared spectrum, 161
- 1,3-propanediol 2-octadecyl, infrared spectrum, 161
- 1,3-propanediol 2-octadecyl, diacetate, infrared spectrum, 161
- 1,3-propanediol 3-octadecyl, ditrifluoroacetate, infrared spectrum, 161
- Ether**, 3-pentyl octadecyl, infrared spectrum, NMR, 161
- synthesis, physical properties, 161
- Ether**, n-propyl octadecyl, infrared spectrum, NMR, 161
- synthesis, physical properties, 161
- Euonymus alatus** (Thunb.) Sieb., diacyl-3-acetin content, 473
- Euonymus europaeus** L., diacyl-3-acetin content, 473
- Euonymus latifolius** (L.) Mill., diacyl-3-acetin content, 473
- Euonymus verrucosus** Scop., diacyl-3-acetin content, 473
- F**
- Fasciola hepatica** L., excretion of lipids, 353
- growth medium, 353
- histochemistry of excretory ducts, 353
- Fasted animals**, lipogenesis in, 60
- See also starvation
- Fat**, bovine milk globule, lipid content, 443
- dog lymph, lipid content, 453
- rat, epididymal tissue, 133
- rat, perirenal tissue, 133
- natural dietary in atherosclerosis, 342
- natural, thermal hydrolysis, 435
- Fat-deficient rats**, erect on fatty acids in phospholipids and triglycerides, 114
- effect on metabolism of unnatural dienoic acids, 122
- Fatty acids**, absorption of methyl esters by rat intestine, 155
- n-alkyl esters of fatty acids, 204
- biosynthesis in fruits, 308
- biosynthesis *in vitro* in *Drosophila melanogaster* larvae, 429
- in bovine milk fat globules, 443
- in bovine milk membranes, 443
- in bovine milk serum, 443
- comp in milk phospholipids, 178
- comp in turnip root lipids, 244
- comp of *Xeranthemum annuum* seed oil, 172
- desaturase-EC 1.3.99? effect of diet, 287
- $\Delta 9,10$ -desaturase-EC 1.3.99? in hen liver, 419
- $\Delta 9,10$ -desaturase-EC 1.3.99? in chicken liver, effect of diet, 461
- $\Delta 9,10$ -desaturase-EC 1.3.99? inhibition by cyclopropane fatty acids, 419
- in dog lymph chylomicron membranes, 453
- in dog lymph fat cores, 453
- in dog lymph serum, 453
- esters isolated by GLC free of contamination, 197
- excreted by *Fasciola hepatica* L., 353
- homologous series of esters related to x-ray long spacings, 204
- incorporation of glucose carbon in dog brain, 195
- isolation from liquid scintillation fluid, 438
- methylation, on an ion exchange resin, 192
- methyl esters from phosphatidyl choline, 479
- from monoene alcohols by reductive ozonolysis, 266
- from phospholipids of yeast extracts, 279
- poly, hydroxy, analysis of isopropylidene derivatives of isomers, 199
- polymorphic forms of esters, 204
- recovery from liquid scintillation fluid, 438
- short chain, quantitative dtm in triglycerides, 473
- synthesis, 225
- synthesis in fasted mice, 60
- TLC, 501
- in Triton hyperlipemia, 212
- Fatty acids, branched-chain**, in milk, 178
- from reductive ring opening of cyclopropane fatty acids, 225
- Fatty acids, chain elongation**, effect of diet, 287
- in chicken liver, effect of diet, 461
- Fatty acids, cyclopropane**, biosynthesis in plant tissues, 308
- geometrical isomers, 225
- positional isomers, 225
- profile of seeds, *Brachychiton acerifolium*, 308
- profile of seeds, *Brachychiton populneum*, 308
- profile of seeds, *Gossypium hirsutum*, 308
- profile of seeds, *Sterculia foetida*, 308
- ring-methylene carbon from methionine, 308
- Fatty acids, cyclopropane**, biosynthesis in plant tissues, 308
- inhibitor of fatty acid $\Delta 9, 10$ -desaturase, 419
- profile of seeds, *Brachychiton acerifolium*, 308
- profile of seeds, *Brachychiton populneum*, 308
- profile of seeds, *Gossypium hirsutum*, 308
- profile of seeds, *Sterculia foetida*, 308
- ring-methylene carbon from methionine, 308
- Fatty acids, epoxy**, in *Xeranthemum annuum* seed oil, 172
- Fatty acids, hydroxy**, column chromatography, 28
- prep, 28
- profile in milk fat, 208
- tri-optimally active from *Chamaepeuce* seed oils, 261
- Fatty acids, positional isomers**, of egg phosphatidyl choline, 149
- of milk serum of phosphatidyl choline, 149
- phosphatidyl choline, determined by phospholipase A, 122
- of safflower seed phosphatidyl choline, 149
- of soybean phosphatidyl choline, 149
- of wheat germ phosphatidyl choline, 149
- Fatty acids, profile**, of bovine milk diglycerides, 443
- of bovine milk phosphatidyl choline, 443
- of bovine milk phosphatidyl ethanolamine, 443
- of bovine milk sphingomyelin, 443
- of bovine milk triglycerides, 443
- of *Brachychiton acerifolium* seed lipids, 316
- of *Brachychiton populneum* seed lipids, 316
- of ceramide aminoethylphosphonate from sea anemone, 55
- of chicken diets, 461
- of chicken liver lipids, effect of diet, 461
- of chicken plasma lipids, effect of diet, 461
- of *Chlorella pyrenoidosa*, 381
- of cholesterol esters in Triton hyperlipemia, 212
- for cod liver lipids, 251
- of corn oil, 287
- of coconut oil, 287
- of dietary fats, 342
- in dietary fats and tissue lipids in rats, 133
- of dog lymph diglycerides, 453
- of dog lymph lysophosphatidyl choline, 453
- of dog lymph phosphatidyl choline, 453
- of dog lymph phosphatidyl ethanolamine, 453
- of dog lymph sphingomyelin, 453
- of dog lymph sterol esters, 453
- of dog lymph triglycerides, 453
- of egg yolk phosphatidyl choline, 217
- in fecal lipid, rats, 155
- of fish oils (hydrogenated), 363
- of intestinal mucosa lipid of rats fed polyunsaturated esters, 155
- in liver, antioxidant deficient rat, 97
- of liver lipids of alloxan-diabetic rats, 284
- of liver phosphatidyl choline, 122
- of menhaden oil triglycerides, 489
- in muscle, antioxidant deficient rat, 80, 100
- of oilseed phospholipids, 88
- of oilseeds, 345
- in phosphatidyl choline in kidney from choline-deficient rats, 239
- in phosphatidyl ethanolamine in kidney from choline-deficient rats, 239
- of phospholipids in hyperlipemia, 212
- in phospholipids and triglycerides of rat epididymal fat pad, 114
- in phospholipids and triglycerides of rat epirenal fat depot, 114
- in phospholipids and triglycerides of rat heart, kidney, lung and pancreas, 114

- in phospholipids and triglycerides of rat intestinal fat depot, 114
- in phospholipids and triglycerides of rat liver lipids, 114
- in *Pullularia pullulans* lipids, 276
- of rat brain, effect of maternal diet, 497
- of rat diets, 497
- from rat intestine, 484
- of rat laboratory chow (Purina), 287
- of rat liver, 287
- of rat liver lipids, effect of diet, 287
- of rat plasma, effect of diet, 287
- of rat stomach contents, 497
- of *Sapindaceae* seed oils, 258
- of seed oils, 258, 371
- of seed oils (hydrogenated), 363
- of sphingomyelin, bovine heart, 21
- of *Sterculia foetida* seed lipids, 316
- in testes, antioxidant deficient rat, 103
- of triglycerides in hyperlipemia, 212
- of whale oil triglycerides, 489
- Fatty acid, unsaturated, arachidonic acid (20:4 ω 6), in antioxidant deficiency in rats, 103**
 - 18:1 ω 9 in cod liver lipids, 251
 - 18:2 ω 6 in cod liver lipids, 251
 - 18:3 ω 3 in cod liver lipids, 251
 - 18:4 ω 3 in cod liver lipids, 251
 - 20:4 ω 6 in cod liver lipids, 251
 - 20:5 ω 3 in cod liver lipids, 251
 - 22:5 ω 3 in cod liver lipids, 251
 - 22:6 ω 3 in cod liver lipids, 251
 - decrease of PUFA in alloxan-diabetic rats, 284
 - dienolic acids, dietary effect in antioxidant deficiency in rats, 89, 97, 103
 - dienolic acids, unnatural, 122
 - flame ionization detector, response of fatty acids, 251
 - geometric configuration, 199
 - GLC, 502
 - hexaenoic acid, dietary effect in antioxidant deficiency in rats, 103
 - linoleic acid (18:2 ω 6), in antioxidant deficiency in rats, 103
 - monoenoic acids, dietary effect in antioxidant deficiency in rats, 89, 97, 103
 - oxidation by permanganate, 199
 - pentaenoic acid, dietary effect in antioxidant deficiency in rats, 103
 - pentaenoic fatty acid (22:5 ω 6), in antioxidant deficiency in rats, 103
 - rat heart, incorporation of 1-¹⁴C-arachidonic acid radio activity, 484
 - tetraenoic acid, dietary effect in antioxidant deficiency in rats, 103
 - TLC of polyunsaturated, 89
 - trienoic acids, dietary effect in antioxidant deficiency in rats, 89, 97, 103
 - in Triton hyperlipemia, 212
- Fatty acid, unsaturated essential, deficiency in rats, 8**
 - deficiency in rats, symptoms, 8
 - deficiency in rats, replacement of di- and tetraenoic acids with mono- and trienoic acids, 8
 - (ω 6) in chick encephalomalacia, 103
- Fatty acid, unsaturated, poly-, biosynthesis, 122**
 - decreased in rat liver in antioxidant deficiency, 97
 - GLC, 502
 - in milk, 178
 - in rat brain, effect of maternal diet, 497
 - reaction with ribonuclease, 83
 - resistant to pancreatic lipase, 489
- Fatty acid, unsaturated, profile, for cod liver lipids, 251**
 - of rat liver phosphatidyl choline, 479
- Fatty acid, unsaturated, PUFA, decrease in rat muscle in antioxidant deficiency, 109**
 - interconversion, 109
 - lower in vitamin E deficient rats, 133
 - in rat muscle decrease in antioxidant deficiency, 89
 - in rat testes decrease in antioxidant deficiency, 103
 - in Triton hyperlipemia, 212
- Feces, excretion of ¹⁴C-cholic acid in suckling rats, 137**
- Feces, human, content of neutral steroids and bile salts, 467**
- Feces, rat, fatty acids in, 155**
- Ferricyanide oxidation, of α -tocopherol, 47**
- Fish, depot fats, fatty acid profile, 251**
- Fish oils, triglycerides, 363**
- Flask design, for incubation under ¹⁸O₂ gas, 33**
- Fluke, liver. See *Fasciola hepatica* L.**
- Formic acid, incorporation into cyclopropane-cyclopropene fatty acids, 308**
- Fructose, effect on lipogenesis in mice, 60**

LIPIDS, VOL. 2, NO. 6

Free radicals, lipid, α -tocopherol oxidation, 47
Freundlich adsorption isotherms. See adsorption isotherms
Fruit. See oil, seeds; oilseeds, or specific name
Fungus. See microorganism

G

- Gadus morhua, cod, liver oil fatty acid profile, 251**
- Galactoglycerides, in alfalfa, 406**
 - infrared spectrum, 406
- Galactolipids, during rat brain development, 76**
- Galactosyl diglyceride, from turnip roots, 244**
- Gangliosides, during rat brain development, 76**
- Gel filtration. Also see chromatography, sephadex**
- Glucose, effect on lipogenesis in mice, 60**
- β -Glucuronidase-EC 3.2.1.3, hydrolysis of α -tocopheronic acid conjugate, 390**
- Glycerides, containing trihydroxy fatty acids from *Chamaepeuce*, 261**
 - fatty acid profile not the same as phospholipids from oilseeds, 88
 - polar fraction in milk fat, 208
 - positional isomers, 149, 217
 - rat liver, incorporation of 1-¹⁴C-arachidonic acid radio activity, 484
 - TLC, 501
- Glycerol, quantitative dtmn in triglycerides, 473**
- Glycerol alkyl ether diesters, in tissue cultures of Ehrlich ascites cells, 193**
- Glycerol ethers, GLC, TLC of isomeric, 161**
 - mono-, isomers chemical and physical properties, 161
 - synthesis, 161
- 1-Glycerol ethers, isopropylidenes, 199**
- Glycerol trimethylsilyl (TMS) ethers, 161**
- Glycerol, 1,2-trimethylsilyl 3-octadecyl ether, infrared spectrum, 161**
- Glycerol, 1,3-trimethylsilyl 2-octadecyl ether, infrared spectrum, 161**
- L- α -Glycerophosphate, incorporation into phosphatidic acid and triglycerides, 14**
 - K_m for triglyceride biosynthesis, 14
- Glycerophosphatides, fatty acid distribution, positional isomers, 149**
- Glycerophosphorylcholine, from phospholipids by yeast extracts, 279**
- L- α -Glyceryl-(2-trimethylammoniumethyl) phosphonate, synthesis and physical properties, 194**
- Glycogen, effect of insulin on synthesis, 60**
- Glycolipids, human brain, 37**
 - spinach leaf, 37
- Glycolysis, in dog brain, 195**
- Glycosyl diglycerides, isolation procedure, 37**
 - spinach leaf, 37
- Goat blood plasma, phospholipid comp, 323**
- Gossypium hirsutum, fatty acid, cyclopropane and cyclopropene, profile of seeds, 308**
 - seed lipids, profile and biosynthesis, 308
- Guinea pig blood plasma, phospholipid comp, 323**
- Gymnosporia royiana M. Laws., diacyl-3-acetin content, 473**

H

- Harpullinase, fatty acid profile, 258**
- Harpullia pendula Planch., fatty acid profile, 258**
- Heart, bovine, lipid extract, 1**
- Heart, rat, fatty acid profile of triglycerides and phospholipids, 114**
 - incorporation of 1-¹⁴C-arachidonic acid radio activity, 484
- Hedeoma drummondii Benth., fatty acid profile, 371**
- Hemorrhagic syndrome, kidney, choline-deficient rats, 239**
- Hepatic. See liver**
- n-Heptadecanoic acid, in *Pullularia pullulans*, 276**
- Hexadecanoic acid, heptadecyl ester, melting point, x-ray long-spacings, 204**
 - hexadecyl ester, melting point, x-ray long-spacings, 204

- octadecyl ester, melting point, x-ray long-spacings, 204
- pentadecyl ester, melting point, x-ray long-spacings, 204
- tetradecyl ester, melting point, x-ray long-spacings, 204
- n-Hexadecanoic acid, in *Pullularia pullulans*, 276
- Hexadecenoic acid, in rat mucosa lipids, 155
- cis-9-Hexadecenoic acid, in synthesis of cyclopropane fatty acids, 225
- Hexose monophosphate pathway, in dog brain, 195
- Hibiscus syriacus, source of 8,9-methyleneheptadecanoic acid, 225
- Homeostatic mechanisms, of tissue lipids comp, 89
- Horse blood plasma, phospholipid comp, 323
- Human, chylomicron in serum of male adults, 183
 - skin surface lipid from scalp, 266
 - tissue cultured cells, 41
- Human, brain, age and lipid comp, 295
- Human carcinoma, of cervix, lipids in tissue cultured HeLa cells, 41
 - of nasopharynx, lipids in tissue cultured KB cells, 41
- Human cell cultures, phospholipid biosynthesis, 424
- Human milk, fatty acid profile of phospholipids, 178
- Human placenta, lipid metabolism, 403
 - phospholipid metabolism, 403
- Human serum, lipoproteins by disc electrophoresis, 282
- Hydnocarpus wightiana, triglyceride profile, 363
- Hydrocarbon, saturated, bovine heart ventricle, 1
 - DEAE column chromatography, 1
- Hydrocarbon, unsaturated, bovine heart ventricle, 1
 - DEAE column chromatography, 1
- Hydrogen iodine value, of unsaturated aldehydes, 280
- Hydrolysis, thermal, of triglycerides, 435
- Hydroperoxidation, of linoleic acid, by lipoxidase, 28
- Hydroperoxide, by lipoxidase, 33
- 9-Hydroperoxyoctadecadienoic acid, methyl ester, lipoxidase product, 28
- 13-Hydroperoxyoctadecadienoic acid, methyl ester, lipoxidase product, 28
- 13-Hydroperoxyoctadeca-9,11-dienoic acid, formation by lipoxidase, from linoleic acid, 432
 - lipoxidase product, 28
 - thermal reaction to form pentane, 432
- 13-Hydrostearic acid, methyl ester, prep, physical properties, 28
- 4-Hydroxydecanoate acid, methyl ester, TMS ether, GLC, 208
- 5-Hydroxydecanoate acid, methyl ester, TMS ether, GLC, 208
- 4-Hydroxydecanoic acid, methyl ester, TMS ether, mass spectral data, major fragmentation ions, 208
- 5-Hydroxydecanoic acid, methyl ester, TMS ether, mass spectral data, major fragmentation ions, 208
 - methyl ester, TMS ether, mass spectral data, major fragmentation ions, 208
- 4-Hydroxydodecanoic acid, methyl ester, TMS ether, mass spectral data, major fragmentation ions, 208
- 5-Hydroxydodecanoic acid, methyl ester, TMS ether, GLC, 208
 - methyl ester, TMS ether, mass spectral data, major fragmentation ions, 208
- 5-Hydroxyheptadecanoic acid, methyl ester, TMS ether, mass spectral data, major fragmentation ions, 208
- 5(?) -Hydroxyheptadecanoic acid, methyl ester, TMS ether, GLC, 208
- 5-Hydroxyhexadecanoic acid, methyl ester, TMS ether,
 - methyl ester, TMS ether, mass spectral data, major fragmentation ions, 208
- Hydroxylation, of double bonds, 199
- Hydroxymethylglutarylcoenzyme A, in cholesterol biosynthesis, 5
- 9 (or 10) -Hydroxymethyl-9-octadecenoic acid, methyl ester, a product from stercularic acid methyl ester with silver nitrate, 316
- 5-Hydroxynonadecanoic acid, methyl ester, TMS ether, mass spectral data, major fragmentation ions, 208
- 5 (?) -Hydroxynonadecanoic acid, methyl ester, TMS ether, GLC, 208
- 5-Hydroxyoctadecanoic acid, methyl ester, TMS ether,
 - methyl ester, TMS ether, mass spectral data, major fragmentation ions, 208
- 9-Hydroxy-trans-10, cis-12-octadecadienoic acid, 33
- 13-Hydroxy-cis-9, trans-11-octadecadienoic acid, 33
- 5-Hydroxyoctanoic acid, methyl ester, TMS ether, GLC, 208
 - methyl ester, TMS ether, mass spectral data, major fragmentation ions, 208
- 5-Hydroxypentadecanoic acid, methyl ester, TMS ether, GLC, 208
 - methyl ester, TMS ether, mass spectral data, major fragmentation ions, 208
- 9-Hydroxystearic acid, nomenclature, 286
- 13-Hydroxystearic acid, methyl ester, from lipoxidase on linoleic acid, 33
- Hydroxystearic acid esters, prep, 28
- 5-Hydroxytetradecanoic acid, methyl ester, TMS ether, GLC, 208
 - methyl ester, TMS ether, mass spectral data, major fragmentation ions, 208
- 4-Hydroxyundecanoate acid, methyl ester, TMS ether, GLC, 208
- 4-Hydroxyundecanoic acid, methyl ester, TMS ether, mass spectral data, major fragmentation ions, 208
- 5-Hydroxyundecanoic acid, methyl ester, TMS ether, GLC, 208
 - methyl ester, TMS ether, mass spectral data, major fragmentation ions, 208
- Hypercholesterolemia, in suckling rat, 137
- Hypocholesterolemia, dietary triglyceride (8:0 and 10:0) effect, 287
- Hyptis decurrens (Blanco) Epling, fatty acid profile, 371
- Hyptis floribunda Briq., fatty acid profile, 371
- Hyptis mutabilis (Rich.) Briq., fatty acid profile, 371
- Hyptis spicigera Lam., fatty acid profile, 371
- Hyptis stellulata Benth., fatty acid profile, 371
- Hyptis suaveolens (L.) Poit., fatty acid profile, 371
- Hyssopus officinalis L., fatty acid profile, 371

- *D-xylo-*, prefix in nomenclature of polyhydroxy fatty acids, 286
- Isomers, diastereo-**, of phytanic acid, 357
- of polyhydroxy fatty acids, 199
- of polyhydroxystearic acids, nomenclature, 286
- of pristanic acid, 357
- Isomers, positional, distributions**, 122
- distribution of fatty acids in egg phosphatidyl choline, 217
- of polyhydroxy fatty acids, 199
- Isopropylidene derivatives**, of polyhydroxy fatty acids, 199

J

- Jurinea antolica* Boiss., pentacyclic triterpenes, 127
- Jurinea consanguinea* DC., fruit, pentacyclic triterpenes, 127

K

- Ketone bodies**, in alloxan-diabetic rats, 231
- Ketostearic acid**, methyl ester, prep, 28
- 13-Ketostearic acid**, methyl ester, from lipoxidase on linoleic acid, 33
- Kidney**, hemorrhagic degeneration in choline-deficient rats, 239
- phospholipid comp in choline-deficient rats, 239
- Kidney, baboon**, acetic acid incorporation into cholesterol, 491
- Kidney, rat**, choline-deficient, 239
- fatty acid profile of triglycerides and phospholipids, 114
- incorporation of 1-¹⁴C-arachidonic acid radio activity, 484
- K_m values**. See enzyme kinetic data
- Koeleruteria bipinnata* Franch., fatty acid profile, 258
- Koeleruteria paniculata* Laxm., fatty acid profile, 258
- Koeleruteriaceae**, fatty acid profile, 258

L

- Labalenic acid**. See 5,6-octadecadienoic acid
- Lactobacillic acid**. See 11,12-methyleneoctadecanoic acid
- Lactone**, precursor in milk fat, 208
- Lallemantia canescens* (L.) Fisch. and Mey., fatty acid profile, 371
- Lallemantia iberica* (M.B.) Fisch. and Mey., fatty acid profile, 371
- Lallemantia royleana* Benth., fatty acid profile, 371
- Lamium amplexicaule* L., fatty acid profile, 371
- Lamium moschatum* Mill., fatty acid profile, 371
- Lamium purpureum* L., fatty acid profile, 371
- Lauraceae**, fatty acid profile; triglyceride profile, 345
- Lauric acid**, in castor seed phospholipids, 88
- Laurus nobilis*, fatty acid profile; triglyceride profile, 345
- Lavandula dentata* L., fatty acid profile, 371
- Lavandula lanata* Boiss., fatty acid profile, 371
- Lavandula latifolia* Medic., fatty acid profile, 371
- Lavandula pendunculata* Cav., fatty acid profile, 371
- Lavandula stoechas* L., fatty acid profile, 371
- Lecithin**, during rat brain development, 76
- sheep erythrocytes, 64
- Leonotis nepetaefolia* (L.) R. Br., fatty acid profile, 371
- Leonurus cardiaca* L., fatty acid profile, 371
- Leonurus sibiricus* L., fatty acid profile, 371
- Lepechinia spicata* Willd., fatty acid profile, 371
- Leucas nutans* Spreng., fatty acid profile, 371
- Lindera benzoin*, fatty acid profile; triglyceride profile, 345
- Lindera praecox*, fatty acid profile; triglyceride profile, 345
- Lindera triloba*, fatty acid profile; triglyceride profile, 345
- Linoleic acid**, isomers after hydroxylation, 286
- Linoleic acid**, isopropylidenes of, GLC, TLC, 199
- Linoleic acid**, autooxidation to form ethane, 432
- in castor seed phospholipids, 88
- decrease in rat muscle phospholipids in antioxidant deficiency, 169

- dietary effect in antioxidant rats, 103
- ester, in rat diet and effect on liver phospholipid fatty acids, 97
- esters, in rat diet and effect on muscle phospholipid fatty acids, 89
- ethyl ester, inhibition of atherogenesis by dietary natural fats, 342
- in galactoglycerides of alfalfa, 406
- hydroperoxidation by lipoxidase, 28
- hydroxylated, geometrical isomers, 199
- incorporation into rat tissues, 114
- isomers after hydroxylation, 286
- isopropylidenes of, GLC, TLC, 199
- lack of effect on fatty acid $\Delta^9,10$ -desaturase, 419
- linolenic acid ratio in maternal rat diet, effect on brain fatty acids, 497
- oxidative desaturation to γ -linolenic acid, 284
- palmityl alcohol ester, preparative procedure, melting point, 437
- in phosphatidyl choline, 149
- in rat muscle phospholipids in antioxidant deficiency, 89
- reaction with ribonuclease, 83
- in Triton hyperlipemia, 212
- in turnip root lipids, 244
- in *Xeranthemum annuum* seed oil, 172
- Linoleic acid, methyl ester**, autooxidation, 28
- hydroperoxidation 33
- in rat diet, 114, 155
- reduction to octadecadienoate isomers, 122
- Linoleic acid, (18:2 ω 6)**, in phosphatidyl choline of rat liver, 122
- cis, cis-Linoleic acid**, isopropylidenes, GLC, 199
- cis, trans-Linoleic acid**, isopropylidenes of, GLC, TLC, 199
- trans- cis-Linoleic acid**, isopropylidenes of, GLC, TLC, 199
- trans, trans-Linoleic acid**, isopropylidenes, GLC, 199
- Linolenic acid**, esters, in rat diet and effect on muscle phospholipid fatty acids, 89
- linoleic acid ratio in maternal rat diet, effect on brain fatty acids, 497
- in turnip roots, 244
- in turnip root lipids, 244
- in *Xeranthemum annuum* seed oil, 172
- Linolenic acid (18:3 ω 3)**, GLC, 502
- α -Linolenic acid**, oxidative desaturation in alloxan-diabetic rats, 284
- γ -Linolenic acid**, from linoleic, 284
- Lipase, pancreatic**, hydrolysis of diacyl-3-acetins, 473
- Lipase, pancreatic-EC 3.1.1.3**, hydrolysis of polyunsaturated fatty acid esters, 489
- hydrolysis of triglycerides, 397
- rates of hydrolysis, 397
- used in preparing diglycerides, 440
- Lipemia, hyper-**, lipid correlation matrix, 212
- lipid correlations, 212
- Lipid extraction**, from alfalfa, 406
- from blood plasma, 323
- from bovine heart ventricle, 1
- from *Chlorella pyrenoidosa*, 381
- from dog lymph, 453
- from dog lymph chylomicrons, 453
- from dog lymph serum, 453
- from dog serum, 212
- from fat under nitrogen, 133
- from feces, 467
- from growth medium for *Fasciola hepatica* L., 353
- from human brain, 295
- from human skin surface, scalp, 266
- from *Jurinea antolica* and *J. consanguinea* fruit, 127
- from milk, bovine, 443
- from milk membranes, bovine, 443
- from milk serum, bovine, 443
- from rabbit liver, 47
- from rat feces, 390
- from rat liver, 47, 390, 479, 484
- from rat liver and serum, fatty acid isolation, 287
- from rat urine, 390
- from seeds, 172, 345, 473
- from sheep erythrocytes, 64
- of tumor lipids, 143
- from turnips, 244
- Lipid free-radicals**, α -tocopherol oxidation, 47
- Lipid metabolism**, effect of dietary triglycerides, 461
- Lipids**, of aging human brain, 295
- beef heart ventricle, 1
- changes in rat brain during development, 76

- comp in turnip roots, 244
- containing free amino groups analysis by trinitrobenzene sulfonic acid, 87
- correlation in experimental endogenous hyperlipemia, 212
- droplets and glycerol ether diesters in Ehrlich ascites cells, 193
- excretion by *Fasciola hepatica* L., 353
- human brain glycolipids, 37
- human skin surface, 266
- of milk, 178
- of newborn dog brain, *in vitro* synthesis, 195
- of sea anemone, 55
- of sheep erythrocytes, 64
- spinach leaf glycolipids, 37
- tissue autooxidation in vitamin E deficient rats, 133
- in tumors, 143
- Lipids, fruit, *Malva nicaensis*, fatty acid biosynthesis, 308**
 - *Malva parviflora*, fatty acid biosynthesis, 308
 - *Malvaviscus arboreus*, fatty acid biosynthesis, 308
- Lipids, oxidizing, reaction with ribonuclease, 83**
- Lipids, seed, *Brachychiton acerifolium*, profile and biosynthesis, 308**
 - *Brachychiton populneum*, profile and biosynthesis, 308
 - *Gossypium hirsutum*, profile and biosynthesis, 308
 - *Sterculia foetida*, profile and biosynthesis, 308
- Lipogenesis, in fasted animals, 60**
 - role of ATP, 60
- Lipomyces. See microorganism**
- Lipoprotein, by disc electrophoresis, 282**
 - human serum, 183
 - infrared quantitation, 183
 - quantitation by elemental analysis, 183
 - recovery calculation, 183
 - very-low-density, triglycerides in alloxan-diabetic rats, 231
- Lipoxidase-EC 1.99.2.1, to form 18-hydroperoxy-9,11-dienoic acid, 482**
 - hydroperoxidation of linoleic acid, 28
 - origin of oxygen incorporated, 33, 442
 - reaction mechanism, 28, 33, 442
- Litchi chinensis, fatty acid profile, 258**
- Lithocholic acid, dtmn in human feces, 467**
- Liver, effect on lipogenesis in mice, 60**
 - electron microscopy in antioxidant deficiency, 97
 - endoplasmic reticulum vacuolization, in antioxidant deficiency, 97
 - endoplasmic reticulum vacuolization, produced by necrogenic diet, 97
 - fatty, in alloxan-diabetic rats, 231
 - hydrolytic activity, rat, 8
 - microsomal supernatant in cholesterol biosynthesis, 137
 - necrogenic diet, torula yeast, reversed by tocopherol, 97
 - perfusion, 231
- Liver, baboon, acetic acid incorporation into cholesterol, 454**
- Liver, chicken, fatty acid $\Delta 9,10$ -desaturase activity, 419**
- Liver, cod, fatty acid profile, 251**
- Liver, rabbit, oxidation of α -tocopherol, 47**
- Liver, rat, cholesterol biosynthesis in suckling rat, 137**
 - cholesterol levels and diet, 287
 - fatty acid profile of, 287
 - fatty acid profile of triglycerides and phospholipids, 114
- incorporation of 1-¹⁴C-archidonate radio activity, 484
- lipid metabolism in experimental diabetes, 231
- oxidation of α -tocopherol, 47
- oxidative desaturation of fatty acids by microsomes, 284
- phosphatidyl choline, fatty acid profile and positional distribution, 122
- phosphatidyl choline, from methionine *in vivo*, 479
- phospholipid comp, effect of antioxidant deficiency, 87
- sterol precursors of cholesterol, 84

- Liver cholesterologenesis, 5**
- Liver fluke. See *Fasciola hepatica* L.**
- Liver-intestine recirculation, of bile, effect on cholesterol biosynthesis in suckling rats, 137**
- Liver lipids, rat, fatty acid profile of, effect of diet, 287**
- Lung, rat, fatty acid profile of triglycerides and phospholipids, 114**
 - incorporation of 1-¹⁴C-archidonate radio activity, 484
- Lupeol, in *Jurinea* fruit, 127**
 - melting point, physical properties, 127
- Lupeol acetate, melting point, physical properties, 127**
- Lycopene, in turnip roots, 244**
- Lycopus asper Greene, fatty acid profile, 371**
- Lycopus europaeus L., fatty acid profile, 371**
- Lymph, dog, effect of diet on lipid comp, 453**
 - lipids of serum and chylomicrons, 453
 - prep of chylomicron and membranes, 453
- Lymphosarcoma, human, lipid, 143**
- Lysocleithin, sheep erythrocytes, 64**
- Lysophosphatidic acid, biosynthesis in intestinal mucosa, 14**
- Lysophosphatidyl choline, fatty acid profile of dog lymph, 453**
 - in determining positional isomers, TLC, 122
 - in dog lymph chylomicron membranes, 453
 - in dog lymph serum, 453
 - infrared spectrum, 353
 - in plasma from various mammals, 323
 - precursor of phosphatidyl choline in human placenta, 403
- Lysophosphatidyl ethanolamine, precursor of phosphatidyl choline in human placenta, 403**
- Lysophospholipase (EC 3.1.1.5.), from *Saccharomyces cerevisiae*, 279**
- Lysosomes, enzymes, 8**
 - in essential fatty acid deficiency, 8
 - membranes, rat liver, 8
 - release of enzymes, 8
 - stability to thermal and osmotic variations, 8
- Lythraceae, fatty acid profile; triglyceride profile, 345**

M

- Malonaldehyde, index of autooxidation of lipids, 133**
 - from polyunsaturated fatty acids, 83
 - reaction with ribonuclease, 83
- Malvalic acid, biosynthesis in plant tissues, 308**
 - inhibitor of fatty acid $\Delta 9,10$ -desaturase, 419
 - methyl ester, products of silver nitrite reaction, 316
 - methyl ester, ring opening on silica gel-silver nitrite TLC plates, 316
- Malva nicaensis, fruit lipids, fatty acid biosynthesis, 308**
- Malva parviflora, fruit lipids, fatty acid biosynthesis, 308**
- Malvaviscus arboreus, fruit lipids, fatty acid biosynthesis, 308**
- Marrubium peregrinum L., fatty acid profile, 371**
- Marrubium pestaloziae Boiss., fatty acid profile, 371**
- Marrubium vulgare L., fatty acid profile, 371**
- Mass spectrometry. See also chromatography, gas-liquid-mass spectrometer**
- Maternal rat diet, effect on brain fatty acids, 497**
- Maturation. See development or growth**
- Maytenus ilicifolia Mart., diacyl-3-acetin content, 473**
- Medicago sativa (alfalfa), galactoglycerides, 496**
- Melissa officinalis L., fatty acid profile, 371**
- Membrane, milk globule membrane lipids, 443**
 - dog lymph chylomicron lipids, 453
 - prep from dog lymph, 453
- Membrane lipids, in antioxidant deficiency, 89**
- Menhaden, oil triglycerides, fatty acid profile, 489**
 - triglyceride profile, 363
- Mentha aquatica L., fatty acid profile, 371**
- Mentha arvensis L., fatty acid profile, 371**
- Mentha longifolia (L.) Huds., fatty acid profile, 371**
- Mentha pulegium L., fatty acid profile, 371**
- Mentha rotundifolia (L.) Huds., fatty acid profile, 371**
- Mentha sylvestris L., fatty acid profile, 371**
- Mentha tomentosa D'Urv., fatty acid profile, 371**
- MER-29. See also 1-p-(β -diethylaminoethoxy) phenol-1-(p-tolyl)-2-(p-chlorophenyl) ethanol or triparanol**
- Metabolism, chemicals affecting lipid, 5**
 - of 7,13-, and 7,13-eicosadienoic acid in rat, 122
 - of 9,15- and 12,15-octadecadienoic acid in rat, 122
 - unsaturated fatty acids in rat intestine, 155
- Methionine, in antioxidant deficiency, effect on peroxidation, 109**
 - in antioxidant deficiency, effect in PUFA, 89
 - methyl carbon incorporation into cyclopropane-cyclopropane fatty acids, 308
 - source of methyl in phosphatidyl choline, rat, 479
- dl-Methionine, effect on antioxidant deficiency in rat, 89, 97, 103**
- N-Methylation, in choline-deficient rats, 239**
- 5,6-Methyleneicosanoic acid, methyl ester, mass spectrum, 225**
- 8,9-Methyleneheptadecanoic acid, from *Hibiscus syriacus*, 225**
 - methyl ester, partial mass spectrum, 225

- 9,10-Methylenehexadecanoic acid, methyl ester, partial mass spectrum, 225
- 9,10-Methylenooctadecanoic acid, methyl ester, partial mass spectrum, 225
- 11,12-Methylenooctadecanoic acid, methyl ester, partial mass spectrum, 225
- 9(or 10)-Methylene-10(or 9)-octadecenoic acid, methyl ester, a product from sterculeic acid methyl ester with silver nitrite, 316
- 9(or 10)-Methylene octade-10-enoic acid, lack of effect on fatty acid $\Delta^9,10$ -desaturase, 419
- 9,10-Methylenetetradecanoic acid, methyl ester, partial mass spectrum, 225
- N-Methyl-N-(3-dimethylamino) propyl-17 β -aminoandrost-5-en-3 β -ol, 20, 25-DAC (SC-12937), cholesterol biosynthesis inhibitor, 5
- Mevalonate, biosynthesis, *in vitro* in liver from suckling rats, 137
- Mevalonic acid, cholesterol biosynthesis, 5
- Mice, fasted, lipogenesis in, 60
- Mice serum, lipoproteins by disc electrophoresis, 282
- Micromeria serpyllifolia (M.B.) Boiss., fatty acid profile, 371
- Microorganism, *Chlorella pyrenoidosa*, culture conditions, 381
- *Chlorella pyrenoidosa*, fatty acid profile, 381
- *Clostridium perfringens* (welchii), phospholipase C (atoxin), 217
- *Lipomyces starkeyi*, lipid comparison with *Pullularia pullulans*, 276
- *Neurospora crassa* strain 34486 for choline assay, 239
- *Pullularia pullulans*, growth medium, 276
- *Pullularia pullulans*, lipid fatty acid profile, 276
- *Saccharomyces cerevisiae*, lipid comparison with *Pullularia pullulans*, 276
- *Saccharomyces cerevisiae* extract, phospholipase activity, 279
- torula yeast, a liver neoprogenic diet, 97
- Microsomes, from cultured human cells, 41
- from cultured human cells, phospholipid biosynthesis, 424
- hamster intestinal mucosa, 14
- oxidative desaturation of fatty acids, 284
- rat kidney, 8
- rat liver, 8
- Milk, bovine, lipids of serum and membranes, 443
- camel, comp of lipids, 178
- cow, comp of lipids, 178
- goat, comp of lipids, 178
- horse, comp of lipids, 178
- human, comp of lipids, 178
- Indian buffalo, comp of lipids, 178
- phosphatidyl choline, fatty acid profile, 178
- phosphatidyl ethanolamines, fatty acid profile, 178
- phosphatidyl serines + phosphatidyl inositols, fatty acid profile, 178
- pig, comp of lipids, 178
- sheep, comp of lipids, 178
- sphingomyelins, fatty acid profile, 178
- Milk fat, hydroxy fatty acid profile, 208
- polar glyceride fraction, 208
- Milk serum, phosphatidyl choline structure, 149
- Mitochondria, from cultured human cells, 41
- from cultured human cells, phospholipid biosynthesis, 424
- Moldavica parviflora (Nutt.) Britt., fatty acid profile, 371
- Molucella laevis L., fatty acid profile, 371
- Molucella spinosa L., fatty acid profile, 371
- Monarda fistulosa L. var. menthaefolia (Graham) Fern., fatty acid profile, 371
- Monarda punctata L., fatty acid profile, 371
- Monardella sp., fatty acid profile, 371
- Monobutyrin, hydrolysis by pancreatic lipase, 397
- Monogalactoglycerides, in alfalfa, 406
- infrared spectrum, 406
- Monoglycerides, TLC, 501
- in triglyceride biosynthesis, intestinal mucosa, 14
- 1-Monoglycerides, isopropylidenes, 199
- Monolayer, compressibility, 21
- of L- α -dipalmitoyl glycerophosphoryl choline, 21
- dipole properties, 21
- effect of calcium on, 21
- gel state, 21
- inclination of the hydrocarbon chains, 21
- liquid state, 21
- solid state, 21
- of sphingomyelin, 21
- surface rheology, 21
- 2-Monopalmitin, K_m for triglyceride biosynthesis, 14
- LIPIDS, VOL. 2, NO. 6
- Mesla punctulata (J. F. Gmel.) Nakai, fatty acid profile, 371
- Mucosa and luminal lipids, rat, fatty acid profile, 155
- Mugil cephalus (mullet), triglyceride profile, 363
- Mullet, triglyceride profile, 363
- Muscle, rat, gastrocnemius, phospholipid fatty acid comp, 89
- phospholipid comp, effect of antioxidant deficiency, 89
- phospholipid fatty acid comp, 109
- quadriceps, phospholipid fatty acid comp, 89
- Muscular dystrophy, nutritional in the rat, contrasted with antioxidant deficiency syndrome, 103
- Mustard seed, erucic acid content, 88
- Myristic acid, in castor seed phospholipids, 88
- myristates of pentacyclic triterpene alcohols, 127
- from oxidative degradation of ceramide aminoethylphosphonate, 55
- in Triton hyperlipemia, 212
- N
- Neomycin, in rat diet, 155
- Neoplastic tissues, lipids in, 143
- See also tumors
- Nepeta cataria L., fatty acid profile, 371
- Nepeta congesta Fisch. and Mey., fatty acid profile, 371
- Nepeta glomerulosa Boiss., fatty acid profile, 371
- Nepeta italica L., fatty acid profile, 371
- Nepeta latifolia DC., fatty acid profile, 371
- Nepeta musinii Spreng., fatty acid profile, 371
- Nepeta nepetella L., fatty acid profile, 371
- Nepeta nuda L. var. albiflora Boiss., fatty acid profile, 371
- Nepeta pannonica L. var. pannonica, fatty acid profile, 371
- Nepeta spicata Benth., fatty acid profile, 371
- Nepeta tmolea Boiss., fatty acid profile, 371
- Nepeta tuberosa L., fatty acid profile, 371
- Neutral lipids, from turnip roots, 244
- Ninhydrin analysis of lipids, difficulty in quantitation, 87
- 9(or 10)-Nitratomethyl-9-octadecenoic acid, methyl ester, a product from sterculeic acid methyl ester with silver nitrite, 316
- NMR. See spectra, nuclear magnetic resonance
- Nomenclature, aliphatic alcohols, short-hand, 266
- polyhydroxy fatty acids, 286
- shorthand system, isomeric glycerol ethers, 161
- Norepinephrine, effect on rat adipose tissue palmitic acid incorporation, 299
- Nuclei, from cultured human cells, 41
- from cultured human cells, phospholipid biosynthesis, 424
- Nucleic acid, dtm procedure, 41
- Nystatin, in rat diet, 155
- O
- Ocimum americanum L., fatty acid profile, 371
- Ocimum basilicum L., fatty acid profile, 371
- Ocimum sanctum L., fatty acid profile, 371
- Ocimum selloi Benth., fatty acid profile, 371
- Octadeca-6,9-dienoic acid, from oleic acid *in vitro*, 284
- 5,6-Octadecadienoic acid, dtm in seed oils, 371
- 9,15-Octadecadienoic acid, synthesis, metabolism in rat, 122
- 12,15-Octadecadienoic acid, synthesis, metabolism in rat, 122
- n-Octadecane, infrared spectrum, 161
- Octadecanoic acid, heptadecyl ester, melting point, x-ray long-spacings, 204
- hexadecyl ester, melting point, x-ray long-spacings, 204
- octadecyl ester, melting point, x-ray long-spacings, 204
- pentadecyl ester, melting point, x-ray long-spacings, 204
- tetradecyl ester, melting point, x-ray long-spacings, 204
- n-Octadecanoic acid, in *Pullularia pullulans*, 276
- Octadecenoic acid, in cod liver phospholipids, 251
- in rat mucosa lipids, 155
- in rat mucosa and luminal lipids, 155
- cis-6-Octadecenoic acid, in synthesis of cyclopropane fatty acids, 225

- cis-9-Octadecenoic acid*, in synthesis of cyclopropane fatty acids, 225
cis-11-Octadecenoic acid, in synthesis of cyclopropane fatty acids, 225
Octadeca-6,9,12,15-tetraenoic acid, from γ -linolenic acid *in vitro*, 284
cis-5, cis-9, cis-12-Octadecatrienoic acid, in *Xeranthemum annuum* seed oil, 172
n-Octadecyl alcohol, infrared spectrum, 161
n-Octyl alcohol, from plastic bag contaminants, 85
2-Octyl-1-cyclopropene-1-heptanoic acid. See *malvalic acid*
2-Octyl-1-cyclopropene-1-octanoic acid. See *sterculic acid*
Oenanthe globulosa L., petroselinic acid content, 339
Oils, coconut, fatty acid profile, 287
 — solvent for cholesterol-Permutit adsorption isotherm, 277
Oils, corn, fatty acid profile of, 287
 — solvent for cholesterol-Permutit adsorption isotherm, 277
Oils, fish, Brevoortia tyrannus (menhaden), triglyceride profile, 363
 — *Enthymus alletteratus* (tuna), triglyceride profile, 363
 — menhaden, oil triglycerides, fatty acid profile, 489
 — menhaden, triglyceride profile, 363
Mugil cephalus (mullet), triglyceride profile, 363
 — mullet, triglyceride profile, 363
 — Filchard oil, triglyceride profile, 363
Sardinops caerulea, Girard, triglyceride profile, 363
 — tuna, triglyceride profile, 363
Oils, marine, polyunsaturated fatty acid profiles, 251
Oils, olive, solvent for cholesterol-Permutit adsorption isotherm, 277
Oils, parinarium, ultra-violet spectrum, 411
Oils, seed, physical properties, 371
Oils, safflower, solvent for cholesterol-Permutit adsorption isotherm, 277
Oils, soybean, hydrogen iodine value, 280
Oils, whale, triglyceride fatty acid profile, 489
 — triglycerides, pancreatic lipase hydrolysis, 489
Oilseeds, Acanthosyrus spinescens, triglyceride profile, 363
 — *Actinodaphne hookeri*, fatty acid profile; triglyceride profile, 345
 — *Actinolema macrolema*, Boiss., petroselinic acid content, 339
 — *Adonis aestivalis* L., diacyl-3-acetin content, 473
 — *Agastache urticifolia* (Benth.) O. Ktze., fatty acid profile, 371
 — *Ajuga chis* (Poir.) Schreb., fatty acid profile, 371
 — *Ajuga iva* Schreb., fatty acid profile, 371
 — *Akebia quinata* Decne, diacyl-3-acetin content, 473
 — *Akebia trifoliata* (Thunb.) Koidz., diacyl-3-acetin content, 473
 — *Amechysta caerulea* L., fatty acid profile, 371
 — *Anisomeles indicia* (L.) O. Ktze., fatty acid profile, 371
 — *Ballota acetabulosa* Benth., fatty acid profile, 371
 — *Ballota hispanica* (L.) Benth., fatty acid profile, 371
 — *Ballota nigra* L. ssp. *ruderalis* (Sw.) Briq., fatty acid profile, 371
 — *Betonica alopecuroides* L., fatty acid profile, 371
 — *Brachychiton acerifolium*, fatty acid profile of seed lipids, 316
 — *Brachychiton acerifolium*, fatty acid, cyclopropane and cyclopentene, profile of seeds, 308
 — *Brachychiton acerifolium*, seed lipids, profile and biosynthesis, 308
 — *Brachychiton populneum*, fatty acid, cyclopropane and cyclopentene, profile of seeds, 308
 — *Brachychiton populneum*, fatty acid profile of seed lipids, 316
 — *Brachychiton populneum*, seed lipids, profile and biosynthesis, 308
 — *Brazoria scutellarioides* Engelm. and Gray., fatty acid profile, 371
 — *Bupleurum croceum* Fenzl., petroselinic acid content, 339
 — *Calamintha clinopodium* Benth., fatty acid profile, 371
 — *Calamintha nepetoides* Jord., fatty acid profile, 371
 — *Cardiospermum canescens* Wall., fatty acid profile, 258
 — *Cardiospermum grandiflorum* Sw., fatty acid profile, 258
 — *Cardiospermum halicababum*, fatty acid profile, 258
 — castor seed phospholipids, fatty acid profile, 88
 — *Celastrus orbiculatus* Thunb., diacyl-3-acetin content, 473
 — *Celastrus scandens* L., diacyl-3-acetin content, 473
 — *Chamaepeuce agra* (Jacq.) DC, 261
 — *Chamaepeuce hispanica* DC, 261
 — *Cinnamomum camphora*, fatty acid profile; triglyceride profile, 345
 — *Cleonia lusitanica* L., fatty acid profile, 371
 — *Cuphea carthagenensis*, fatty acid profile; triglyceride profile, 345
 — *Cuphea hookeriana*, fatty acid profile; triglyceride profile, 345
 — *Cuphea ignea*, fatty acid profile; triglyceride profile, 345
 — *Cuphea lanceolata*, fatty acid profile; triglyceride profile, 345
 — *Cuphea lavia*, fatty acid profile; triglyceride profile, 345
 — *Cuphea painteri*, fatty acid profile; triglyceride profile, 345
 — *Daucus guttatus* Sibth. & Sm., petroselinic acid content, 339
 — *Dodonea*, fatty acid profile, 258
 — *Dodonea lanceolata* F. Meull., fatty acid profile, 258
 — *Dodonea viucosa* Jacq., fatty acid profile, 258
 — *Dracocephalum moldavica* L., fatty acid profile, 371
 — *Ducroisia anethifolia* Boiss., petroselinic acid content, 339
 — *Elakoltzia patrini* Garcke, fatty acid profile, 371
 — *Elakoltzia splendens* Nak., fatty acid profile, 371
 — *Eremostachys laciniata* (L.) Bunge, fatty acid profile, 371
 — *Eremostachys speciosa* Rupr., fatty acid profile, 371
 — *Eremostachys vicaryi* Benth., fatty acid profile, 371
 — *Eryngium eburneum* Decne., petroselinic acid content, 339
 — *Euonymus alatus* (Thunb.) Sieb., diacyl-3-acetin content, 473
 — *Euonymus europaeus* L., diacyl-3-acetin content, 473
 — *Euonymus latifolius* (L.) Mill., diacyl-3-acetin content, 473
 — *Euonymus verrucosus* Scop., diacyl-3-acetin content, 473
 — *Fasciola hepatica* L., excretion of lipids, 353
 — *Fasciola hepatica* L., growth medium, 353
 — *Fasciola hepatica* L., histochemistry of excretory ducts, 353
 — *Gossypium hirsutum*, fatty acid, cyclopropane and cyclopentene, profile of seeds, 308
 — *Gossypium hirsutum*, seed lipids, profile and biosynthesis, 308
 — *Gymnosporia royleana* M. Laws., diacyl-3-acetin content, 473
 — *Harpullia pendula* Planch., fatty acid profile, 258
 — *Harpullia*, fatty acid profile, 258
 — *Hedeoma drummondii* Benth., fatty acid profile, 371
 — *Hydnocarpus wightiana*, triglyceride profile, 363
 — *Hyptis decurrens* (Blanco) Epling, fatty acid profile, 371
 — *Hyptis floribunda* Briq., fatty acid profile, 371
 — *Hyptis mutabilis* (Rich.) Briq., fatty acid profile, 371
 — *Hyptis spicigera* Lam., fatty acid profile, 371
 — *Hyptis stellulata* Benth., fatty acid profile, 371
 — *Hyptis suaveolens* (L.) Poit., fatty acid profile, 371
 — *Hyssopus officinalis* L., fatty acid profile, 371
 — *Jurinea antolica* Boiss., pentacyclic triterpenes, 127
 — *Jurinea consanguinea* D.C., fruit, pentacyclic triterpenes, 127
 — *Koeleruteria bipinnata* Franch., fatty acid profile, 258
 — *Koeleruteria paniculata* Laxim., fatty acid profile, 258
 — *Koeleruteriaceae*, fatty acid profile, 258
 — *Lallemantia canescens* (L.) Fisch. and Mey., fatty acid profile, 371
 — *Lallemantia iberica* (M. B.) Fisch. and Mey., fatty acid profile, 371
 — *Lallemantia royleana* Benth., fatty acid profile, 371
 — *Lamium amplexicaule* L., fatty acid profile, 371
 — *Lamium moschatum* Mill., fatty acid profile, 371
 — *Lamium purpureum* L., fatty acid profile, 371
 — *Lauraceae*, fatty acid profile; triglyceride profile, 345
 — *Laurus nobilis*, fatty acid profile; triglyceride profile, 345
 — *Lavandula dentata* L., fatty acid profile, 371
 — *Lavandula lanata* Boiss., fatty acid profile, 371
 — *Lavandula latifolia* Medic., fatty acid profile, 371
 — *Lavandula pendunculata* Cav., fatty acid profile, 371
 — *Lavandula stoechas* L., fatty acid profile, 371
 — *Leonotis nepetifolia* (L.) R. Br., fatty acid profile, 371
 — *Leonurus cardiaca* L., fatty acid profile, 371
 — *Leonurus sibiricus* L., fatty acid profile, 371
 — *Lepechinia spicata* Willd., fatty acid profile, 371
 — *Leuca nufana*, Spreng., fatty acid profile, 371
 — *Lindera benzoin*, fatty acid profile; triglyceride profile, 345
 — *Lindera praecox*, fatty acid profile; triglyceride profile, 345

- *Lindera triloba*, fatty acid profile; triglyceride profile, 345
- *Litchi chinensis*, fatty acid profile, 258
- *Lycopus asper* Greene, fatty acid profile, 371
- *Lycopus europaeus* L., fatty acid profile, 371
- *Lythraceae*, fatty acid profile; triglyceride profile, 345
- *Marrubium peregrinum* L., fatty acid profile, 371
- *Marrubium pestalozzae* Boiss., fatty acid profile, 371
- *Marrubium vulgare* L., fatty acid profile, 371
- *Maytenus illicifolia* Mart., diacyl-3-acetin content, 473
- *Melissa officinalis* L., fatty acid profile, 371
- *Mentha aquatica* L., fatty acid profile, 371
- *Mentha arvensis* L., fatty acid profile, 371
- *Mentha longifolia* (L.) Huds., fatty acid profile, 371
- *Mentha pulegium* L., fatty acid profile, 371
- *Mentha rotundifolia* (L.) Huds., fatty acid profile, 371
- *Mentha sylvestris* L., fatty acid profile, 371
- *Mentha tomentosa* D'Urv., fatty acid profile, 371
- *Micromeria scryphifolia* (M.B.) Boiss., fatty acid profile, 371
- *Moldavica parviflora* (Nutt.) Britt., fatty acid profile, 371
- *Molucella laevis* L., fatty acid profile, 371
- *Molucella spinosa* L., fatty acid profile, 371
- *Monarda fistulosa* L. var. *menthaefolia* (Graham) Fern., fatty acid profile, 371
- *Monarda punctata* L., fatty acid profile, 371
- *Monardella* sp., fatty acid profile, 371
- *Mosla punctulata* (J. F. Gmel.) Nakai, fatty acid profile, 371
- mustard seed, erucic acid content, 88
- *Nepeta cataria* L., fatty acid profile, 371
- *Nepeta congesta* Fisch. and Mey., fatty acid profile, 371
- *Nepeta glomerulosa* Boiss., fatty acid profile, 371
- *Nepeta italica* L., fatty acid profile, 371
- *Nepeta latifolia* DC., fatty acid profile, 371
- *Nepeta mussinii* Spreng., fatty acid profile, 371
- *Nepeta nepetella* L., fatty acid profile, 371
- *Nepeta ruda* L. var. *albiflora* Boiss., fatty acid profile, 371
- *Nepeta pannonica* L. var. *pannonica*, fatty acid profile, 371
- *Nepeta spicata* Benth., fatty acid profile, 371
- *Nepeta tmolea* Boiss., fatty acid profile, 371
- *Nepeta tuberosa* L., fatty acid profile, 371
- *Ocimum americanum* L., fatty acid profile, 371
- *Ocimum basilicum* L., fatty acid profile, 371
- *Ocimum sanctum* L., fatty acid profile, 371
- *Ocimum scitoides* Benth., fatty acid profile, 371
- *Oenanthe globulosa* L., petroselinic acid content, 339
- *Origanum vulgare* L., fatty acid profile, 371
- *Otostegia limbata* (Benth.) Boiss., fatty acid profile, 371
- *Paulinia cupana* H. B. and K., fatty acid profile, 258
- *Pauliniaeae*, fatty acid profile, 258
- *Perilla frutescens* (L.) Britt., fatty acid profile, 371
- *Petroselinum crispum* (Mill.) Mansf., petroselinic acid content, 339
- *Phlomis armeniaca* Willd., fatty acid profile, 371
- *Phlomis austra-anatolica* Hub. Mur., fatty acid profile, 371
- *Phlomis crinita* Cav., fatty acid profile, 371
- *Phlomis fruticosa* L., fatty acid profile, 371
- *Phlomis herba-venti* L., fatty acid profile, 371
- *Phlomis lycia* D. Don, fatty acid profile, 371
- *Phlomis purpurea* L., fatty acid profile, 371
- *Phlomis rigida*, Labill., fatty acid profile, 371
- *Phyostegia virginiana* (L.) Benth., fatty acid profile, 371
- *Plectranthus inflexus* (Thunb.) Vahl ex Benth., 371
- *Pogostemon parviflorus* Benth., fatty acid profile, 371
- *Prasium majus* L., fatty acid profile, 371
- *Prunella asiatica* Nak., fatty acid profile, 371
- *Prunella vulgaris* L., fatty acid profile, 371
- *Prunella vulgaris* L. var. *lanceolata* (Bart.) Fern., fatty acid profile, 371
- *Pycnanthemum muticum* Pers., fatty acid profile, 371
- rape seed, erucic acid content, 88
- *Rosmarinus officinalis* L., fatty acid profile, 371
- safflower seed, phosphatidyl choline structure, 149
- *Salvia acetabulosa* L. var. *simplicifolia* Boiss., fatty acid profile, 371
- *Salvia aegyptiaca* L., fatty acid profile, 371
- *Salvia aethiops* L., fatty acid profile, 371
- *Salvia amplexicaulis* Lam., fatty acid profile, 371
- *Salvia apiana* Jeps., fatty acid profile, 371
- *Salvia bicolor* Desf., fatty acid profile, 371
- *Salvia brachyantha* (Bordz.) Pobedim., fatty acid profile, 371
- *Salvia bracteata* Russ., fatty acid profile, 371
- *Salvia cardiaca* Benth., fatty acid profile, 371
- *Salvia ceratophylla* L., fatty acid profile, 371
- *Salvia coccinea* Juss. ex Murr., fatty acid profile, 371
- *Salvia columbariae* Benth., fatty acid profile, 371
- *Salvia cryptantha* Monthr. and Auch., fatty acid profile, 371
- *Salvia euphratica* Monthr. and Auch., fatty acid profile, 371
- *Salvia farinacea* Benth., fatty acid profile, 371
- *Salvia glutinosa* L., fatty acid profile, 371
- *Salvia grandiflora* Etling., fatty acid profile, 371
- *Salvia hispanica* L., fatty acid profile, 371
- *Salvia horminum* L., fatty acid profile, 371
- *Salvia hydrangea* DC. ex Benth., fatty acid profile, 371
- *Salvia judaica* Boiss., fatty acid profile, 371
- *Salvia lanigera* Poir., fatty acid profile, 371
- *Salvia lavendulaefolia* Vahl., fatty acid profile, 371
- *Salvia longispicata* Mart. and Gal., fatty acid profile, 371
- *Salvia lyrata* L., fatty acid profile, 371
- *Salvia mexicana* L., fatty acid profile, 371
- *Salvia montbretti* Benth., fatty acid profile, 371
- *Salvia moercreftiana* Wall. ex Benth., fatty acid profile, 371
- *Salvia nemorosa* L., fatty acid profile, 371
- *Salvia officinalis* L., fatty acid profile, 371
- *Salvia polybeia* R. Br., fatty acid profile, 371
- *Salvia polystachya* Ortt., fatty acid profile, 371
- *Salvia pratensis* L., fatty acid profile, 371
- *Salvia reflexa* Hornem., fatty acid profile, 371
- *Salvia ringens* Sibth. and Sm., fatty acid profile, 371
- *Salvia rooseae* Sm., fatty acid profile, 371
- *Salvia rugosa* Dryand. ex Ait., fatty acid profile, 371
- *Salvia sclarea* L., fatty acid profile, 371
- *Salvia similata* Haussk., fatty acid profile, 371
- *Salvia sonomensis* Greene, fatty acid profile, 371
- *Salvia suffruticosa* Monthr. and Auch., fatty acid profile, 371
- *Salvia sylvatica* L., fatty acid profile, 371
- *Salvia syriaca* L., fatty acid profile, 371
- *Salvia tchikatcheffii* (Fisch. and Mey.) Boiss., fatty acid profile, 371
- *Salvia texana* (Sheele) Torr., fatty acid profile, 371
- *Salvia triloba* L. f., fatty acid profile, 371
- *Salvia valentina* Vahl., fatty acid profile, 371
- *Salvia verbenaca* L., fatty acid profile, 371
- *Salvia verticillata* L., fatty acid profile, 371
- *Sapindaceae*, fatty acid profile, 258
- *Sapindus drummondii* H. and A., fatty acid profile, 258
- *Sapindus mukorossi* Gaertn., fatty acid profile, 258
- *Sapindus saponaria* L., fatty acid profile, 258
- *Sassafras albidum*, fatty acid profile; triglyceride profile, 345
- *Satureja coreana* (Levl.) Nakai, fatty acid profile, 371
- *Satureja hortensis* L., fatty acid profile, 371
- *Satureja thymra* L., fatty acid profile, 371
- *Scutellaria columbae* Ait., fatty acid profile, 371
- *Scutellaria condensata* Rech. f., fatty acid profile, 371
- *Scutellaria drummondii* Benth., fatty acid profile, 371
- *Scutellaria multicaulis* Boiss., fatty acid profile, 371
- *Sideritis incana* Benth., fatty acid profile, 371
- *Sideritis hirsuta* L., fatty acid profile, 371
- *Sideritis lagasacna* Willk., fatty acid profile, 371
- *Sideritis leucantha* Cav., fatty acid profile, 371
- *Sideritis linearifolia* Lam., fatty acid profile, 371
- *Sideritis montana* L. var. *comosa* Boiss., fatty acid profile, 371
- *Sideritis taurica* Willd., fatty acid profile, 371
- *Sideritis tragoriganum* Lag., fatty acid profile, 371
- *Sorbus aucuparia* L., diacyl-3-acetin content, 473
- *Sorbus maugeoti* Godr. and Soy-Will., diacyl-3-acetin content, 473
- soybean, phosphatidyl choline structure, 149
- *Stachys arvensis* L., fatty acid profile, 371
- *Stachys cretica* Sibth. and Sm., fatty acid profile, 371
- *Stachys germanica* L., fatty acid profile, 371
- *Stachys hirta* L., fatty acid profile, 371
- *Stachys milanii* Petrov., fatty acid profile, 371
- *Stachys olympica* Poir., fatty acid profile, 371
- *Stachys sylvatica* L., fatty acid profile, 371
- *Stachys thirkei* C. Koch., fatty acid profile, 371
- *Stachys viticina* Boiss., fatty acid profile, 371
- *Sterculia foetida*, fatty acid, cyclopropane and cyclopropene, profile of seeds, 308
- *Sterculia foetida*, fatty acid profile of seed lipids, 316

- *Sterculia foetida*, seed lipids, profile and biosynthesis, 308
- *Sterculia foetida*, triglycerides profile, 363
- *Tetradlea coulteri* Gray, fatty acid profile, 371
- *Teucrium almeriense* C. E. Hubb. and Sandw., fatty acid profile, 371
- *Teucrium capitatum* L., fatty acid profile, 371
- *Teucrium chamaedrys* L., fatty acid profile, 371
- *Teucrium creticum* L., fatty acid profile, 371
- *Teucrium cubense* Jacq. ssp. *laevigatum* (Vahl) McClint. and Epl., fatty acid profile, 371
- *Teucrium depressum* Small, fatty acid profile, 371
- *Teucrium expannum* Pau, fatty acid profile, 371
- *Teucrium flavum* L., fatty acid profile, 371
- *Teucrium gnaphalodes* Vahl, fatty acid profile, 371
- *Teucrium polium* L., fatty acid profile, 371
- *Teucrium pseudochamaepitys* L., fatty acid profile, 371
- *Teucrium scoroides* Schreb., fatty acid profile, 371
- *Teucrium scorodonia* L., fatty acid profile, 371
- *Thymra spicata* L., fatty acid profile, 371
- *Thymus capitatus* (L.) Hoffm. and Link, fatty acid profile, 371
- *Thymus chaubardii* Boiss. and Heldr. ex Boiss., fatty acid profile, 371
- *Thymus vulgaris* L., fatty acid profile, 371
- *Thymus zygis* L., fatty acid profile, 371
- *Ulmaceae*, fatty acid profile; triglyceride profile, 345
- *Ulmus americana*, fatty acid profile; triglyceride profile, 345
- *Ulmus pumila*, fatty acid profile; triglyceride profile, 345
- *Umbellularia californica*, fatty acid profile; triglyceride profile, 345
- *Wiedemannia orientalis* Fisch. and Mey., fatty acid profile, 371
- *Xanthoceras sorbifolia* Bunge, fatty acid profile, 258
- *Xeranthemum annuum*, fatty acid comp, 172
- *Zelkova serrata*, fatty acid profile; triglyceride profile, 345
- *Ziziphora capitata* L., fatty acid profile, 371
- *Ziziphora tenuior* L., fatty acid profile, 371
- Oleic acid**, in castor seed phospholipids, 88
- conversion to PUFA in rat antioxidant deficiency, 109
- hydrogen iodine value, 280
- isopropylidenes of, GLC, TLC, 199
- in milk, 178
- oxidative desaturation in alloxan-diabetic rats, 284
- in phosphatidyl choline, 149
- in phosphatidyl choline of rat liver, 122
- in rat phospholipids and triglycerides, 114
- in Triton hyperlipemia, 212
- in turnip root lipids, 244
- in *Xeranthemum annuum* seed oil, 172
- Oleic acid (18:1)**, GLC, 502
- Oleyl acetate**, preparative procedure, melting point, 437
- Olive oil**, solvent for cholesterol-Permutit adsorption isotherm, 277
- Optical rotatory dispersion**, of methyl 9-D-hydroxyoctadecanoate, 172
- of methyl 9,10-epoxyoctadecanoate, 172
- of methyl *cis*-9-L, 10-L, epoxy-*cis*-12-octadecenoate, 172
- of methyl *cis*-12-D, 13-D-epoxy-*cis*-9-octadecenoate, 172
- of methyl 12-D, 13-D-epoxyoctadecanoate, 172
- of methyl 12-D-hydroxyoctadecanoate, 172
- of 1,9-D-octadecanediol, 172
- of 1,9- and 1,10-octadecanediol, 172
- of 1,12-D-octadecanediol, 172
- of 1,12-D- and 1,13-D-octadecanediols, 172
- Origanum vulgare** L., fatty acid profile, 371
- Otostegia limbata** (Benth.) Boiss., fatty acid profile, 371
- Oxidation**, of linoleic acid and ribonuclease, 83
- Oxidative chain cleavage**, 225
- Oxidative desaturation**, of fatty acids, 284
- P**
- Palmitic acid**, in castor seed phospholipids, 88
- incorporation into di- and triglycerides by rat adipose tissue, 299
- incorporation into phosphatidic acid and triglycerides, 14
- in milk, 178
- oxidative desaturation in alloxan-diabetic rats, 284
- palmitates of penicillate triesterene alcohols, 127
- in triglyceride synthesis in alloxan-diabetic rats, 231
- in Triton hyperlipemia, 212
- in turnip root lipids, 244
- in *Xeranthemum annuum* seed oil, 2:172
- Palmitin-2, 3-dibutyryl**, hydrolysis by pancreatic lipase, 397
- synthesis, 397
- Palmitoleic acid**, cholesterol ester, preparative procedure, melting point, 437
- conversion to PUFA in rat antioxidant deficiency, 109
- isopropylidenes of, GLC, TLC, 199
- from palmitic acid *in vitro*, 284
- in phosphatidyl choline of rat liver, 122
- in rat phospholipids and triglycerides, 114
- in Triton hyperlipemia, 212
- Palmitoyl-Coenzyme A**, in biosynthesis of triglyceride, 14
- hydrolase-EC 3.1.2.2. low activity in hamster intestinal mucosa microsomes and supernatant, 14
- inhibitors of acylation in intestinal mucosa, 14
- Pancreas**, rat, fatty acid profile of triglycerides and phospholipids, 114
- Pancreatic lipase-EC 3.1.1.3.**, acetyl diacylglycerol structure dtmn, 217
- acetyl 1,2-diacylglyceride structure dtmn, 149
- hydrolysis of polyunsaturated fatty acid esters, 489
- phosphatidyl choline structure dtmn, 149, 217
- See lipase, pancreatic
- Parinaric acid**, polarography, 411
- triglycerides, ultra-violet spectrum, 411
- See 9,11,13,15-octadecatetraenoic acid
- Parinarium oil**, ultra-violet spectrum, 411
- Paulinia cupana** H.B. and K., fatty acid profile, 258
- Paulinias**, fatty acid profile, 258
- Pelargonaldehyde**, double bond titration by borohydride, 250
- n-Pentadecanoic**, in *Pullularia pullulana*, 276
- 2-Pentenal**, in vitamin E deficient rats, 133
- Pentane**, from 13-hydroperoxyoctadeca-9,11-dienoic acid by thermal decomposition, 432
- Perilla frutescens** (L.) Britt., fatty acid profile, 371
- Permutit**, adsorption of cholesterol, 277
- Peroxidation**, of 7-dehydrocholesterol, 5
- kinetics in rat muscle in antioxidant deficiency, 109
- of PUFA in antioxidant deficiency, 109
- Peroxides**, absent in vitamin E deficient rats, 133
- lipid, α -tocopherol oxidation, 47
- 5 α , 8 α -Peroxide**, of 7-dehydrocholesterol, 5
- Petroselinic acid**, dtmn by microreactor chromatography, 339
- isopropylidenes of, GLC, TLC, 199
- ozonolysis, 339
- in seed oils, 339
- See *cis*-6-octadecenoic acid
- Petroselinum crispum** (Mill.) Mansf., petroselinic acid content, 339
- Phenazine methosulfate**, effect on lipid synthesis *in vitro*, 195
- 1-p-(β -diethylaminoethoxy) Phenol-1-(p-tolyl)-2-(p-chlorophenol) ethanol**, triparanol or Mer-29, cholesterol biosynthesis inhibitor, 5
- Phlomis lycia** D. Don, fatty acid profile, 371
- Phloionolic acid**, See (+)-*threo*-9,10-18-Trihydroxyoctadecanoic acid
- Phlomis armeniaca** Willd., fatty acid profile, 371
- Phlomis austro-anatolica** Hub. Mur., fatty acid profile, 371
- Phlomis crinita** Cav., fatty acid profile, 371
- Phlomis fruticosa** L., fatty acid profile, 371
- Phlomis herba-venti** L., fatty acid profile, 371
- Phlomis purpurea** L., fatty acid profile, 371
- Phlomis rigida** Labill., fatty acid profile, 371
- Phosphatase**, acid, subcellular distribution in essential fatty acid deficiency in rats, 8
- L- α -Phosphatidate phosphohydrolase-EC 3.1.3.4.**, in hamster intestinal mucosa supernatant fraction, 14
- Phosphatidic acid**, biosynthesis in intestinal mucosa, 14
- from egg phosphatidyl choline, 14
- incorporation of glucose carbon in dog brain, 195
- sheep erythrocytes, 64
- Phosphatidyl choline**, biosynthesis, in choline-deficient rats, 239
- bovine heart ventricle, 1
- in bovine milk membranes, 443
- in bovine milk serum, 443
- calcium interaction with monolayer, 21
- in cultured human cells, 41
- DEAE column chromatography, 1
- in dog lymph chylomicron membranes, 453
- in dog lymph serum, 453
- egg, structure of, 149
- excreted by *Fasciola hepatica* L., 353
- fatty acid profile in bovine milk, 443
- fatty acid profile of dog lymph, 453
- in human brain, age, 295
- hydrolysis by yeast extract, 279
- incorporation of glucose carbon in dog brain, 195
- infrared spectrum, 323

- in *in vivo* incorporation of ^{14}C -methyl from methionine, rat liver, 479
- milk, fatty acid profile, 178
- milk serum, structure of, 149
- molecular species dtmn., 217
- monolayer properties, 21
- phosphorus incorporation in cultured human cells, 424
- in plasma from various mammals, 323
- rat liver, fat-deficient rat, 122
- rat liver, fatty acid profile and positional distribution, 122
- safflower seed, structure of, 149
- in sea anemone phospholipids, 55
- separation according to extent of unsaturation, 479
- soybean, structure of, 149
- structure dtmn., 149, 217
- synthesis in human placenta, 403
- TLC, preparative, 479
- in turnip roots, 244
- wheat germ, structure of, 149
- Phosphatidyl ethanolamine**, analysis by trinitrobenzene sulfonic acid, 87
 - bovine heart ventricle, 1
 - in bovine milk membranes, 443
 - in bovine milk serum, 443
 - in choline-deficient rat kidney, 239
 - in cultured human cells, 41
 - DEAE column chromatography, 1
 - in dog lymph chylomicron membranes, 453
 - in dog lymph serum, 453
 - during rat brain development, 76
 - excreted by *Fasciola hepatica* L., 353
 - fatty acid profile in bovine milk, 443
 - fatty acid profile of dog lymph, 453
 - in human brain, age, 295
 - incorporation of glucose carbon in dog brain, 195
 - infrared spectrum, 64, 323
 - milk, fatty acid profile, 178
 - phosphorus incorporation in cultured human cells, 424
 - in plasma from various mammals, 323
 - in sea anemone phospholipids, 55
 - sheep erythrocytes, 64
 - synthesis in human placenta, 403
 - in turnip roots, 244
- Phosphatidyl glycerol**, bovine heart ventricle, 1
 - DEAE column chromatography, 1
 - in turnip roots, 244
- Phosphatidyl inositol**, bovine heart ventricle, 1
 - in cultured human cells, 41
 - DEAE column chromatography, 1
 - during rat brain development, 76
 - infrared spectrum, 323
 - milk, fatty acid profile, 178
 - phosphorus incorporation in cultured human cells, 424
 - in plasma from various mammals, 323
 - sheep erythrocytes, 64
 - in turnip roots, 244
- Phosphatidyl inositol + sphingomyelin**, incorporation of glucose carbon in dog brain, 195
- Phosphatidyl serine**, analysis by trinitrobenzene sulfonic acid, 87
 - bovine heart ventricle, 1
 - in cultured human cells, 41
 - DEAE column chromatography, 1
 - during rat brain development, 76
 - in human brain, age, 295
 - incorporation of glucose carbon in dog brain, 195
 - milk, fatty acid profile, 178
 - phosphorus incorporation in cultured human cells, 424
 - in sea anemone phospholipids, 55
 - sheep erythrocytes, 64
- Phospholipase**, rat intestinal mucosa, different from other tissue phospholipase, 303
 - rat tissues, trypsin activation, 303
 - of yeast, 279
- Phospholipase A**, use in determining positional isomers of phosphatidyl choline, 122
- Phospholipase A-EC 3.1.1.4.**, from *Saccharomyces cerevisiae*, 279
 - in human placenta, 403
 - fatty acid isomers of phospholipids, dtmn., 178
- Phospholipase B-EC 3.1.1.5.**, rat intestinal mucosa, different from other tissue phospholipase B, 303
 - rat tissues, trypsin activation, 303
- Phospholipase C-EC 3.1.4.3.**, not present in *Saccharomyces cerevisiae*, 279
 - use in phosphatidyl choline structure dtmn., 217
- Phospholipase-EC 3.1.4.3.**, use in phosphatidyl choline structure dtmn., 149
- Phospholipids**, biosynthesis, in human tissue cultures, 424
 - of bovine blood plasma, 323
 - in bovine milk components, 443
 - of cat blood plasma, 323
 - in cultured human cells, 41
 - of dog blood plasma, 323
 - in dog lymph components, 453
 - dtmn procedure, 41
 - egg yolk, 217
 - fatty acid comp, 239
 - fatty acid comp of rat muscle in antioxidant deficiency, 89
 - fatty acid profile in cod liver, 251
 - fatty acid profiles in fat deficient rat, 114
 - fatty acid profile in hyperlipemia, 212
 - fatty acid profile not the same as glycerides from oil-seeds, 88
 - of goat blood plasma, 323
 - of guinea pig blood plasma, 323
 - of horse blood plasma, 323
 - human brain, 37
 - incorporation of glucose carbon in dog brain, 195
 - incorporation of palmitate by perfused rat liver, 231
 - kidney, comp in choline-deficient rats, 239
 - in livers of alloxan-diabetic rats, 231
 - metabolism in human placenta, 403
 - milk, fatty acid profile, 178
 - phosphorus incorporation, in human tissue cultures, 424
 - of pig blood plasma, 323
 - PUFA, chain elongation in antioxidant deficiency, 89
 - of rabbit blood plasma, 323
 - of rat blood plasma, 323
 - rat liver, incorporation of 1-14-C-arachidonic acid radio activity, 484
 - of sea anemone, 55
 - of sheep blood plasma, 323
 - sheep erythrocytes, 64
 - TLC, 501
 - in triglyceride biosynthesis, intestinal mucosa, 14
 - in turnip roots, 244
- Phosphonic acids**, in sea anemone phospholipids, 55
- Phosphonolipids**, 194
 - in sea anemone, 55
 - structural dtmn., 55
- Phthalic acid**, esters, as contaminate from plastic bags, 85
- Physostegia virginiana** (L.) Benth., fatty acid profile, 371
- Phytanic acid**, diastereoisomers dtmn by GLC, 357
 - dtmn by GLC, 357
- Pig blood plasma**, phospholipid comp, 323
- Pigments**, fluorescent of the lipofuscin or ceroid type, in antioxidant deficiency, 89, 97
- Pilchard** oil, triglyceride profile, 363
- Placenta**, human, acyl Co A: acyl GPE transferase, 403
 - phospholipase A activity, 403
 - phospholipid metabolism, 403
- Plasma**, blood, phospholipids of various mammals, 323
 - bovine, blood, TLC, 501
 - rat, fatty acid profile of, effect of diet, 287
- Plasmalogen**, dtmn by 2,4-dinitrophenylhydrazinophosphoric acid reagent, 334
 - ethanolamine, ethanolamine incorporation in rat brain, 329
 - sheep erythrocytes, 64
- Plectranthus inflexus** (Thunb.) Vahl ex Benth., fatty acid profile, 371
- Pogostemon parviflorus** Benth., fatty acid profile, 371
- Polarography**, conjugated unsaturated lipids, 411
- Polarography, half-wave potentials**, all-*trans*-retinal, 411
 - all-*trans*-retinol, 411
 - all-*trans*-retinyl acetate, 411
 - β -carotene, 411
 - 13-*cis*-retinol, 411
 - eleostearic acid, 411
 - ergosterol, 411
 - parinaric acid, 411
 - Vitamin D₂, 411
 - Vitamin D₃, 411
- Polyacrylamide gel**, use in disc electrophoresis of lipoprotein, 282
- Polyglycerol phosphatide**, in cultured human cells, 41
 - phosphorus incorporation in cultured human cells, 424
- Polyunsaturated fatty acid**. See fatty acid, unsaturated
- Potassium**, concentration in sheep erythrocytes, 64
- Potato**, lipids, TLC, 244
- Prasium majus** L., fatty acid profile, 371
- Pristanic acid**, diastereoisomers dtmn by GLC, 357
 - dtmn by GLC, 357
- Propanal**, absence in vitamin E deficient rats, 133
- 1,2-Propanediol 3-octadecyl ether diacetate**, infrared spectrum, 161
- 1,2-Propanediol 3-octadecyl ether ditrifluoroacetate**, infrared spectrum, 161
- 1,3-Propanediol 2-octadecyl ether**, infrared spectrum, 161
- 1,3-Propanediol 2-octadecyl ether diacetate**, infrared spectrum, 161

1,3-Propanediol 3-octadecyl ether ditrifluoroacetate, infrared spectrum, 161
 Prostaglandin E₁, hydroxylation analogous to lipoxidase oxidation, 33
 Protein, in bovine milk, 443
 — in cultured human cells, 41
 — in dog lymph chylomicron membranes, 453
 — dtmn in cultured human cells, 424
 — dtmn procedure, 41
Prunella asiatica Nak., fatty acid profile, 371
Prunella vulgaris L., fatty acid profile, 371
Prunella vulgaris L. var. *lanceolata* (Bart.) Fern., fatty acid profile, 371
 PUFA. See fatty acid, unsaturated
 Pullularia. See microorganism
Pycnanthemum muticum Pers., fatty acid profile, 371
 Pyridoxine deficiency, rat, arachidonic acid metabolism, 484

R

Rabbit blood plasma, phospholipid comp, 323
 Rabbit liver, α -tocopherol oxidation, 47
 Rabbit serum, lipoproteins by disc electrophoresis, 282
 Radio assay, GLC peaks, 484
 — from TLC of phosphatidyl choline, 479
 Rape seed, erucic acid content, 88
 Rat, alloxan-diabetic, liver lipid metabolism, 231
 — arachidonic acid metabolism in pyridoxine deficiency, 484
 — choline-deficient, 239
 — effect of diet on cholesterol synthesis, 287
 — essential fatty acid deficiency, 8
 — laboratory chow (Purina), fatty acid profile of, 287
 — maternal diet, effect on brain fatty acids, 497
 — metabolism of α -tocopherol hydroquinone, 390
 — metabolism of α -tocopherol quinone, 390
 — suckling, cholesterol biosynthesis in liver, 137
 — suckling, cholic acid turnover, 137
 — suckling, hypercholesterolemia, 137
 — unsaturated fatty acid metabolism, 155
 — urinary excretion of α -tocopheronic acid conjugate, 390
 — vitamin E deficient, 133
 — Walker 256 carcinoma, diacylglycerol ethers in human lymphosarcoma, 143
 Rat adipose tissue, palmitic acid incorporation into di- and triglycerides, 299
 Rat blood plasma, phospholipid comp, 323
 Rat brain, ethanolamine incorporation into lipid ethers, 329
 — incorporation of 1-¹⁴C-arachidonic acid radio activity, 484
 Rat heart, incorporation of 1-¹⁴C-arachidonic acid radio activity, 484
 Rat intestinal mucosa, phospholipases, 303
 Rat intestine, fatty acid profile, 484
 — incorporation of 1-¹⁴C-arachidonic acid radio activity, 484
 Rat kidney, incorporation of 1-¹⁴C-arachidonic acid radio activity, 484
 Rat liver, incorporation of 1-¹⁴C-arachidonic acid radio activity, 484
 — phosphatidyl choline, fatty acid profile and positional distribution, 122
 — phosphatidyl choline, from methionine *in vivo*, 479
 — sterol precursors of cholesterol, 84
 — α -tocopherol oxidation, 47
 Rat liver homogenates, cholesterol biosynthesis and inhibitors, 5
 Rat lung, incorporation of 1-¹⁴C-arachidonic acid radio activity, 484
 Rat muscle, phospholipid fatty acid comp, and effect of antioxidant deficiency, 89, 109
 Rat serum, lipoproteins by disc electrophoresis, 282
 Rat testes, incorporation of 1-¹⁴C-arachidonic acid radio activity, 484
 — neutral lipid fatty acid comp and effect of antioxidant deficiency, 103
 — phospholipid fatty acid comp and effect of antioxidant deficiency, 103
 Rat tissues, fatty acid profiles in fat deficient rat, 114
 — phospholipases, 303
 Δ^7 -Reductase, 7-dehydrocholesterol- Δ^7 -reductase, rat liver, 5
 Δ^24 -Reductase, 24-dehydrosterol- Δ^24 -reductase, rat liver, 5
 Reductive ring opening, 225
 Renal (epirenal) fat depot, rat, fatty acid profile of triglycerides and phospholipids, 114
 Renal (perirenal) fat, in vitamin E deficient rats, 133
 Renol. See kidney

all-trans-Retinal, polarography, 411
 all-trans-Retinel, polarography, 411
 13-cis-Retinel, polarography, 411
 all-trans-Retinyol acetate, polarography, 411
 Ribonuclease, reaction with linoleic acid, 83
 — reaction with malonaldehyde, 83
 — reaction with polyunsaturated fatty acids, 83
 Ribonucleic acid, in cultured human cells, 41
 Ricinoleic acid, hydroxylated, geometrical isomers, 199
 — isopropylidenes of, GLC, TLC, 199
 Ricinoleic acid, absent from castor seed phospholipids, 88
 — hydroxylated, geometrical isomers, 199
 — isopropylidenes of, GLC, TLC, 199
Rosmarinus officinalis L., fatty acid profile, 371
 Ruminant milk, fatty acid profile of phospholipids, 178

S

Saccharomyces. See microorganism
 Safflower oil, solvent for cholesterol-Permutit adsorption isotherm, 277
 Safflower seed, phosphatidyl choline structure, 149
Salvia acetabulosa L. var. *simplicifolia* Boiss., fatty acid profile, 371
Salvia aegyptiaca L., fatty acid profile, 371
Salvia aethiops L., fatty acid profile, 371
Salvia amplexicaulis Lam., fatty acid profile, 371
Salvia apiana Jeps., fatty acid profile, 371
Salvia bicolor Desf., fatty acid profile, 371
Salvia brachyantha (Horda.) Pobedim., fatty acid profile, 371
Salvia bracteata Russ., fatty acid profile, 371
Salvia carduacea Benth., fatty acid profile, 371
Salvia ceratophylla L., fatty acid profile, 371
Salvia coccinea Juss. ex Murr., fatty acid profile, 371
Salvia columbariae Benth., fatty acid profile, 371
Salvia cryptantha Monthr., and Auch., fatty acid profile, 371
Salvia euphratica Monthr. and Auch., fatty acid profile, 371
Salvia farinacea Benth., fatty acid profile, 371
Salvia glutinosa L., fatty acid profile, 371
Salvia grandiflora Edling., fatty acid profile, 371
Salvia hispanica L., fatty acid profile, 371
Salvia horminum L., fatty acid profile, 371
Salvia hydrangea DC. ex Benth., fatty acid profile, 371
Salvia judaica Boiss., fatty acid profile, 371
Salvia lanigera Poir., fatty acid profile, 371
Salvia lavandulaefolia Vahl, fatty acid profile, 371
Salvia longispicata Mart. and Gal., fatty acid profile, 371
Salvia lyrata L., fatty acid profile, 371
Salvia mexicana L., fatty acid profile, 371
Salvia montbretii Benth., fatty acid profile, 371
Salvia mucrofratiana Wall. ex Benth., fatty acid profile, 371
Salvia nemorosa L., fatty acid profile, 371
Salvia officinalis L., fatty acid profile, 371
Salvia plebeia R. Br., fatty acid profile, 371
Salvia polystachya Ort., fatty acid profile, 371
Salvia pratensis L., fatty acid profile, 371
Salvia reflexa Hornem., fatty acid profile, 371
Salvia ringens Sibth. and Sm., fatty acid profile, 371
Salvia rosaeifolia Sm., fatty acid profile, 371
Salvia rugosa Dryand. ex Ait., fatty acid profile, 371
Salvia sclarea L., fatty acid profile, 371
Salvia similata Hausk., fatty acid profile, 371
Salvia sonomensis Greene, fatty acid profile, 371
Salvia suffruticosa Monthr. and Auch., fatty acid profile, 371
Salvia sylvestris L., fatty acid profile, 371
Salvia syriaca L., fatty acid profile, 371
Salvia tchihatcheffii (Fisch. and Mey.) Boiss., fatty acid profile, 371
Salvia texana (Sheele) Torr., fatty acid profile, 371
Salvia triloba L. f., fatty acid profile, 371
Salvia valentina Vahl, fatty acid profile, 371
Salvia verbenaca L., fatty acid profile, 371
Salvia verticillata L., fatty acid profile, 371
 Sapindaceae, fatty acid profile, 258
Sapindus drummondii H. and A., fatty acid profile, 258
Sapindus mukorossi Gaertn., fatty acid profile, 258
Sapindus saponaria L., fatty acid profile, 258
Sardinops caerulea, Girard, triglyceride profile, 263
Sassafras albidum, fatty acid profile; triglyceride profile, 345
Satureja coreana (Levl.) Nakai, fatty acid profile, 371
Satureja hortenensis L., fatty acid profile, 371
Satureja thymbrala L., fatty acid profile, 371
 SC-11952. See 20 α -(2-dimethylaminoethyl) amino-5 α -pregnan-3 β -ol dihydrochloride

- SC-12937. See N-methyl-N-(3-dimethylamino) propyl-17 β -amino- α -rost-5-en-3 β -ol
- Scalp, human, surface lipids, 266
- Scintillation fluid, recovery of fatty acid from, 438
- Scutellaria columnae All., fatty acid profile, 371
- Scutellaria condensata Rech. f., fatty acid profile, 371
- Scutellaria drummondii Benth., fatty acid profile, 371
- Scutellaria multicaulis Boiss., fatty acid profile, 371
- Sea anemone, *Anthopleura elegantissima*, phospholipids, 55
- Sebum, human, 266
- Selachyl alcohol, infrared spectrum, 143
- Selenium, in antioxidant deficiency effect on peroxidation, 109
- in antioxidant deficiency effect on, PUFA, 89, 97, 103
- Sephadex. See chromatography, sephadex
- Serine, analysis by trinitrobenzene sulfonic acid, 87
- from oxidative degradation of ceramide aminoethyl-phosphonate, 55
- Serum, lymph lipids, 453
- milk lipids, 443
- Serum albumin, disc electrophoresis, 282
- Serum lipids, in Triton hyperlipemia, 212
- Serum lipoprotein, by disc electrophoresis, 282
- Sheep, Hampshire, erythrocyte lipids, 64
- Sheep, blood plasma, phospholipid comp, 323
- Sheep milk, fatty acid profile of phospholipids, 178
- Sideritis hirsuta L., fatty acid profile, 371
- Sideritis incana Benth., fatty acid profile, 371
- Sideritis lagascana Willk., fatty acid profile, 371
- Sideritis leucantha Cav., fatty acid profile, 371
- Sideritis linearifolia Lam., fatty acid profile, 371
- Sideritis montana L. var. comosa Boiss., fatty acid profile, 371
- Sideritis taurica Willd., fatty acid profile, 371
- Sideritis tragoriganum Lag., fatty acid profile, 371
- β -Sitosterol, in turnip roots, 244
- β -Sitosterol glucoside, in turnip root, 244
- Sodium, concentration in sheep erythrocytes, 64
- Sodium selenite, effect in antioxidant deficiency in rat, 89, 97, 103
- Soil samples, stored in standard soil bags, contaminated by phthalic acid esters, 55
- Sorbus aucuparia L., diacyl-3-acetin content, 473
- Sorbus maugeoti Godr. and Soy-Will., diacyl-3-acetin content, 473
- Soybean, phosphatidyl choline, reference in analysis of turnip root lipids, 244
- phosphatidyl choline structure, 149
- Soybean lipoxidase, 28
- Soybean oil, hydrogen iodine value, 280
- Spectra, infrared, absorption bands of triglycerides, 183
- cholesterol, 64
- didecyl phthalate, 85, 198
- digalactoglyceride, 406
- of 18:0-2 glycerol ether diacetate, 161
- of 18:0-1 and 18:0-2 isomeric glycerol ethers, 161
- of long-chain synthetic mono- and diunsaturated 1- and 2-glycerol monoethers, 161
- lysophosphatidyl choline, 323
- monogalactoglyceride, 406
- n-octadecane, 161
- of cis-5, cis-9, cis-12-octadecatrienoic acid methyl ester, 172
- n-octadecyl alcohol, 161
- 3-pentyl octadecyl ether, 161
- phosphatidyl choline, 323
- phosphatidyl ethanolamine, 64, 323
- phosphatidyl inositol, 323
- 1,2-propanediol 3-octadecyl ether, 161
- 1,2-propanediol 3-octadecyl ether, diacetate, 161
- 1,2-propanediol 3-octadecyl ether ditrifluoroacetate, 161
- 1,3-propanediol 2-octadecyl ether, 161
- 1,3-propanediol 2-octadecyl ether diacetate, 161
- 1,3-propanediol 2-octadecyl ether ditrifluoroacetate, 161
- n-propyl octadecyl ether, 161
- selachyl alcohol, 143
- sheep erythrocyte lipids, 64
- sphingomyelin, 64, 323
- 1,2-trimethylalyl 3-octadecyl glycerol ether, 161
- 1,3-trimethylalyl 2-octadecyl glycerol ether, 161
- Spectra, mass, branched acid esters from methyl 5,6-methyleneecanoate, 225
- branched acid esters from methyl 8,9-methyleneheptadecanoate, 225
- branched acid esters from methyl 9,10-methylenehexadecanoate, 225
- branched acid esters from methyl 9,10-methyleneoctadecanoate, 225
- branched acid esters from methyl 9,10-methylenetradecanoate, 225
- branched acid esters from methyl 11,12-methyleneoctadecanoate, 225
- branched fatty acid, origins of peaks, 225
- 4-hydroxydecanoic acid, 208
- 5-hydroxydecanoic acid, 208
- of 5-hydroxydecanoic acid, methyl ester, TMS ether, 208
- 4-hydroxydodecanoic acid, 208
- 5-hydroxydodecanoic acid, 208
- 5-hydroxyheptadecanoic acid, 208
- 5-hydroxyhexadecanoic acid, 208
- 5-hydroxynonadecanoic acid, 208
- 5-hydroxyoctadecanoic acid, 208
- 5-hydroxyoctanoic acid, 208
- 5-hydroxypentadecanoic acid, 208
- 9-hydroxystearic acid, methyl ester, 28
- 13-hydroxystearic acid, methyl ester, 28, 33
- 5-hydroxytetradecanoic acid, 208
- 4-hydroxyundecanoic acid, 208
- 5-hydroxyundecanoic acid, 208
- Spectra, nuclear magnetic resonance, diacyl-3-acetins, 473
- of 18:1-1 glycerol ether diacetate, 161
- of isomeric glycerol ether, 161
- of cis-5, cis-9, cis-12-octadecatrienoic acid methyl ester, 172
- of 3-pentyl octadecyl ether, 161
- of n-propyl octadecyl ether, 161
- threo-9,10,18-trihydroxy-cis-12-octadecenoic acid, methyl ester, 261
- threo-9,10,18-trihydroxy-cis-12-octadecenoic acid, methyl ester, after D₂O exchange, 261
- Spectra, optical rotatory dispersion, diacyl-3-acetins, 473
- (S)-1,2-distearin-3-acetin, 473
- Spectra, ultra-violet, maxima for carotenoids, 244
- parinaric acid containing triglycerides, 411
- Spectra, x-ray, of n-alkyl esters of fatty acids, 204
- long-spacings, of n-alkyl esters of fatty acids, 204
- Sperm, content, in antioxidant deficiency in rat, 103
- effect of selenium and methionine, 103
- Spermatogenesis, in antioxidant deficiency, 103
- Sphingolipids, during rat brain development, 76
- Sphingomyelin, bovine heart fatty acid profile, 21
- bovine heart ventricle, 1
- in bovine milk membranes, 443
- in bovine milk serum, 443
- calcium interaction with monolayer, 21
- in cultured human cells, 41
- DEAE column chromatography, 1
- dipole estimation, 21
- in dog lymph chylomicron membranes, 453
- in dog lymph serum, 453
- electrophoresis of aqueous dispersion, 21
- ethanolamine incorporation in rat brain, 329
- excreted by *Fasciola hepatica* L., 353
- fatty acid profile in bovine milk, 443
- fatty acid profile of dog lymph, 453
- in human brain, age, 295
- infrared spectrum, 64, 323
- milk, fatty acid profile, 178
- monolayer properties, 21
- phosphorus incorporation in cultured human cells, 424
- in plasma from various mammals, 323
- sheep erythrocytes, 64
- Sphingosine, in sea anemone phosphonolipids, 55
- Spinach leaf, glycolipids, 37
- Spleen, baboon, acetic acid incorporation into cholesterol, 494
- Stachys arvensis L., fatty acid profile, 371
- Stachys cretica Sibth. and Sm., fatty acid profile, 371
- Stachys germanica L., fatty acid profile, 371
- Stachys hirta L., fatty acid profile, 371
- Stachys olympica Poir., fatty acid profile, 371
- Stachys milanii Petrov., fatty acid profile, 371
- Stachys sylvatica L., fatty acid profile, 371
- Stachys thirkei C. Koch, fatty acid profile, 371
- Stachys viticina Boiss., fatty acid profile, 371
- Starvation. See also fasted animals
- Stearic acid, in castor seed phospholipids, 88
- desaturation in chicken liver, effect of diet, 461
- Δ 9, 10-desaturation by hen liver, 419
- methyl ester, from lipoxidase on linoleic acid, 33
- methyl ester, prep, 28
- oxidative desaturation in alloxan-diabetic rats, 284
- in rat mucosa and luminal lipids, 155
- in rat phospholipids and triglycerides, 114
- in Triton hyperlipemia, 212
- in *Xeranthemum annuum* seed oil, 172
- Stearic acid (18:0), GLC, 502
- Sterculia foetida, fatty acid, cyclopropane and cyclopropene, profile of seeds, 308
- fatty acid profile of seed lipids, 316
- seed lipids, profile and biosynthesis, 308
- triglyceride profile, 363
- Sterculic acid, biosynthesis in plant tissues, 308
- inhibitor of fatty acid Δ 9,10-desaturase, 419
- methyl ester, products of silver nitrite reaction, 316

- methyl ester, ring opening on silica gel-silver nitrite TLC plates, 316
- Sterols**, acetic acid incorporation into baboon brain, 494
 - acetylation, 72
 - biosynthesis by rat liver, 5
 - neutral, excretion by humans, 467
 - in turnip roots, 244
- Sterol acetates**, column chromatography, 72
 - GLC, 72
- Sterol esters**, fatty acid profile of dog lymph, 453
 - in human skin surface lipids, 266
 - in turnip roots, 244
- Sterol glucosides**, in potatoes, 244
 - in turnip roots, 244
- Stigmasterol glucoside**, in turnip root, 244
- Subcellular alterations**, in antioxidant deficiency, 97
- Subcellular distribution**, of acid phosphatase in rat liver and kidney, 8
 - of aryl sulfatase in rat liver and kidney, 8
- Subcellular fractions**, chemical comp of human cells, 41
 - of rat liver, prep, 8
 - tissue culture, prep, 41
- Sulfatase**, aryl, subcellular distribution in essential fatty acid deficiency in rats, 8
- Sulfatides**, during rat brain development, 76
 - human brain, 37
 - isolation procedure, 37
 - See also cerebroside sulfate
- Sulfolipid**, isolation procedure, 37
 - spinach leaf, 37
- Sulfur amino acids**, in antioxidant deficiency, effect on PUFA, 89, 103
- Supernatant fraction**, intestinal mucosa, required for triglyceride biosynthesis, 14
- Surface potential**, monolayer, L- α -dipalmitoyl glycerophosphoryl choline, 21
 - effect of calcium, 21
 - sphingomyelin and dihydrosphingomyelin, 21
- Surface pressure**, monolayer, L- α -dipalmitoyl glycerophosphoryl choline, 21
 - effect of calcium, 21
 - sphingomyelin and dihydrosphingomyelin, 21

T

- Taraxasterol**, in *Jurinea* fruit, 127
 - melting point, physical properties, 127
- ψ -Taraxasterol**, in *Jurinea* fruit, 127
 - melting point, physical properties, 127
- Taraxasterol acetate**, melting point, physical properties, 127
- ψ -Taraxasterol acetate**, melting point, physical properties, 127
- Teeth**, incisor depigmentation, in vitamin E deficient rats, 133
- Testes**, rat, incorporation of 1-¹⁴C-arachidonic acid radioactivity, 484
 - phospholipid comp, effect of antioxidant deficiency, 103
- Testicular degeneration**, due to dietary antioxidant deficiency, 103
- Tetraclea coalteri** Gray, fatty acid profile, 371
- Tetracos-14-en-1-ol**, in human skin lipids, 266
- Tetracycline**, in rat diet, 155
- Tetradecanoic acid**, heptadecyl ester, melting point, x-ray long-spacings, 204
 - hexadecyl ester, melting point, x-ray long-spacings, 204
 - octadecyl ester, melting point, x-ray long-spacings, 204
 - pentadecyl ester, melting point, x-ray long-spacings, 204
 - tetradecyl ester, melting point, x-ray long-spacings, 204
- n-Tetradecanoic acid**, in *Pullularia pullulans*, 276
- cis-9-Tetradecenoic acid**, in synthesis of cyclopropane fatty acids, 225
- Tetrahydroxystearic acids**, diastereoisomers, 199
- erythro-9,10-erythro-12,13-Tetrahydroxystearic acid**, nomenclature, 286
- threo-9,10-threo-12,13-Tetrahydroxystearic acid**, nomenclature, 286
- 9,10,12,13-Tetrahydroxystearic acids**, melting points of isomers, 286
 - methyl esters, isomers, 199
- 13-Tetrahydroxystearic acids**, nomenclature, 286
- 2,5,7,8-Tetramethyl-2-(4,8,12-trimethyltridecyl)-6-chromanol**. See α -tocopherol
- 3,7,11,15-Tetramethylhexadecanoic acid**, in cod liver oils, 251
 - See phytanic acid
- 1-(2,5-dihydroxy-3,4,4-trimethylphenyl)-3,7,11,15-Tetramethyl-3-hexadecanol**. See α -tocopheryl hydroquinone
- 2,6,10,14-Tetramethylpentadecanoic acid**, in cod liver oil, 251
 - See pristanic acid
- Teucrium almeriense** C. E. Hubb. and Sandw., fatty acid profile, 371
- Teucrium capitatum** L., fatty acid profile, 371
- Teucrium chamaedrys** L., fatty acid profile, 371
- Teucrium creticum** L., fatty acid profile, 371
- Teucrium cubense** Jacq. ssp. *laevigatum* (Vahl) McClint. and Epl., fatty acid profile, 371
- Teucrium depressum** Small, fatty acid profile, 371
- Teucrium expansum** Pau, fatty acid profile, 371
- Teucrium flavum** L., fatty acid profile, 371
- Teucrium gnaphalodes** Vahl, fatty acid profile, 371
- Teucrium polium** L., fatty acid profile, 371
- Teucrium pseudochamaepitys** L., fatty acid profile, 371
- Teucrium scorodonia** L., fatty acid profile, 371
- Teucrium scordoides** Schreb., fatty acid profile, 371
- Thermal hydrolysis**, of triglycerides, 435
- Thermal reaction**, of 13-hydroperoxyoctadeca-9,11-dienoic acid, 432
- Thoracic lymph**, dog, lipids of serum and chylomicrons, 453
- Thymra apicata** L., fatty acid profile, 371
- Thymus capitatus** (L.) Hoffm. and Link, fatty acid profile, 371
- Thymus chaubardii** Boiss. and Heldr. ex Boiss., fatty acid profile, 371
- Thymus vulgaris** L., fatty acid profile, 371
- Thymus zygis** L., fatty acid profile, 371
- Tissue culture**, glycerol ether diesters in Ehrlich ascites cells, 193
 - growth medium, human, 424
- Tissue cultured cells**, HeLa, 41
 - HeLa, phospholipid biosynthesis, 424
 - human heart, 41
 - human heart, phospholipid biosynthesis, 424
 - human liver, phospholipid biosynthesis, 424
 - KB, 41
 - KB, phospholipid biosynthesis, 424
 - liver tissue, 41
 - subcellular particle prep, 424
- Tissue culture media**, 41
- α -Tocopherol**, oxidation by ferricyanide, 47
 - oxidation in rabbit liver, 47
 - oxidation in rat liver, 47
- α -Tocopherol deficiency**. See also antioxidant deficiency
- α -Tocopherol dimer**, from rat liver, 47
- α -Tocopherol trimer**, from rabbit liver, 47
 - from rat liver, 47
- α -Tocopheronic acid**, conjugate, hydrolysis by β -glucuronidase, 390
 - urinary excretion of conjugate, rat, 390
- α -Tocopheryl hydroquinone**, metabolism in rats, 390
- Tocopheryl-p-quinone**, from rabbit liver, 47
 - from rat liver, 47
- α -Tocopheryl quinone**, metabolism in rats, 390
- Tridecanol**, internal standard, 217
 - in seed oil, 345
- Trifluoroacetic acid**, esters of glycerol ethers, 161
- Triglycerides**, analytical method for glycerol, 473
 - analytical method for short chain fatty acids, 473
 - biosynthesis by α -glycerophosphate pathway, 14
 - biosynthesis in hamster intestinal mucosa, 14
 - bovine heart ventricle, 1
 - in bovine milk fat globules, 443
 - in bovine milk membranes, 443
 - in bovine milk serum, 443
 - comp of seed fats (oilseeds), 345
 - containing trihydroxy fatty acids, from *Chamaepeuce*, 261
 - DEAE column chromatography, 1
 - in diet and lipid metabolism, 287
 - in dog lymph chylomicron membranes, 453
 - in dog lymph fat cores, 453
 - in dog lymph serum, 453
 - effect on lipid metabolism, chicken, 461
 - excreted by *Fasciola hepatica* L., 353
 - fatty acid profile in bovine milk, 443
 - fatty acid profile of cod liver, 251
 - fatty acid profile of dog lymph, 453
 - fatty acid profile in fat deficient rat, 114
 - fatty acid profile in hyperlipemia, 212
 - fish oils, 363
 - hydrolysis, thermal, 435
 - incorporation of glucose carbon in dog brain, 195

- incorporation of palmitate by perfused rat liver, 231
- infrared absorption bands, 183
- *in vitro* biosynthesis stimulation, 14
- menhaden oil, fatty acid profile, 489
- norepinephrine effect on palmitic acid incorporation, 299
- output by perfused rat liver, 231
- palmitic acid incorporation by rat adipose tissue, 299
- polarography, 411
- rat liver lipid metabolism, 287
- seed oils, 363
- stereospecific analysis, 440
- synthesis and utilization, alloxan-diabetic rat liver, 231
- TLC, 601
- in turnip roots, 244
- whale oil, fatty acid profile, 489
- of cholesterol, atherosclerosis model, 277
- three-9,10,18-Trihydroxyoctadecanoic acid, methyl ester, NMR spectrum, 261
- (+)-three-9,10,18-Trihydroxyoctadecanoic acid, from *Chamaepeuce*, 261
- (+)-three-9,10,18-Trihydroxy-cis-12-octadecenoic acid, from *Chamaepeuce*, 261
- Trihydroxystearic acid, methyl esters, 199
- erythro-9, 10-erythro-10, 12-Trihydroxystearic acid, methyl esters, GLC, TLC, 199
- erythro-9, 10-three-10, 12-Trihydroxystearic acid, methyl esters, GLC, TLC, 199
- three-9, 10-erythro-10, 12-Trihydroxystearic acid, methyl esters, GLC, TLC, 199
- three-9, 10-three-10, 12-Trihydroxystearic acid, methyl esters, GLC, TLC, 199
- 9,10,12-Trihydroxystearic acids, melting points of isomers, 286
- methyl esters, isomers, 199
- nomenclature, 286
- Triaurin, in seed oil, 345
- (3-hydroxy-3,7,11,15-tetramethylhexadecyl)-Trimethyl-p-benzoquinone. See α -tocopheryl quinone
- 4,8,12-Trimethyltridecanoic acid, in cod liver oils, 251
- Trinitrobenzene sulfonic acid, analysis of lipids containing free amino groups, 87
- Triolein, hydrolysis by pancreatic lipase, 397
- Triparanol, cholesterol biosynthesis inhibitor, 5, 84
- See also 1-p-(β -diethylaminoethoxy) phenol-1-(p-tolyl)-2-(p-chlorophenol) ethanol
- Triterpenes, pentacyclic, of fruit of *Jurinea anatolica* Boiss. and *J. consanguinea* D. C., 127
- Triton, induced hyperlipemia, 212
- Trypsin, phospholipase activation, 303
- Tumor, human lymphosarcoma, diacylglycerol ethers in human lymphosarcoma, 143
- lipids in, 143
- Walker 256 carcinoma diacylglycerol ethers in human lymphosarcoma, 143
- Tuna, triglyceride profile, 363
- Turnip root, lipids, 244

U

- Umbellularia californica*, fatty acid profile; triglyceride profile, 345
- Ulmaceae*, fatty acid profile; triglyceride profile, 345
- Ulmus americana*, fatty acid profile; triglyceride profile, 345
- Ulmus pumila*, fatty acid profile; triglyceride profile, 245
- Ultracentrifuge, isolation of serum chylomicron, 183
- prep of salt gradient, 183
- Unsaturation, dtmn in oils in presence of aldehydes, 280
- Unsaturated lipids, polarography, 411
- Urea adducts, of aliphatic alcohol acetates, 266
- Urine creatine and creatinine, in antioxidant deficiency, 89

V

- cis- and trans-Vaccenic acid, isopropylidenes of, GLC, TLC, 199
- Ventricle, heart. See heart
- Vitamin B₆. See pyridoxine
- Vitamin D₂, polarography, 411
- Vitamin D₃, polarography, 411
- Vitamin E, antioxidant *in vivo*, 47
- See α -tocopherol
- Vitamin E deficient rat, autooxidation of tissue lipids, 133

W

- Water, in human brain, age, 295
- Wax alcohols, of human skin surface lipids, 266
- Wax alcohol acetates, in human skin surface lipids, 266
- Wax esters, of human skin lipids, 266
- preparative procedure, 437
- Whale, oil triglyceride fatty acid profile, 489
- oil triglycerides, pancreatic lipase hydrolysis, 489
- Wheat germ, phosphatidyl choline structure, 149
- Wiedemannia orientalis Fisch. and Mey., fatty acid profile, 371

X - Y - Z

- Xanthoceras sorbifolia* Bunge, fatty acid profile, 258
- X-irradiation of rats, effect on phospholipases, 303
- X-ray diffraction, of n-alkyl esters of fatty acids, 204
- See also spectra, x-ray
- Zelkova serrata*, fatty acid profile; triglyceride profile, 345
- Ziziphora capitata* L., fatty acid profile, 371
- Ziziphora tenuior* L., fatty acid profile, 371



